
Traditional Chinese medicine — *Lycium barbarum* and *Lycium chinense* fruit

Médecine traditionnelle chinoise — Baie de goji (baie de Lycium barbarum et de Lycium chinense)

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 23193:2020](https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020)

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>



iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 23193:2020

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2020

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

	Page
Foreword.....	v
Introduction.....	vi
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions.....	1
4 Descriptions.....	2
5 Requirements.....	4
5.1 Morphological features.....	4
5.1.1 Appearance.....	4
5.1.2 Colour.....	4
5.1.3 Dimensions.....	4
5.1.4 Fracture.....	4
5.1.5 Odour.....	4
5.2 Microscopic characteristics.....	4
5.3 Moisture.....	5
5.4 Total ash.....	5
5.5 Acid-insoluble ash.....	5
5.6 Water-soluble extractives.....	5
5.7 Thin-layer chromatogram (TLC) identification.....	6
5.8 Content of polysaccharide.....	6
5.9 Content of marker compound.....	6
5.10 Heavy metals.....	6
5.11 Pesticide residues.....	6
5.12 Sulfur dioxide residues.....	6
6 Sampling.....	6
7 Test methods.....	7
7.1 Macroscopic identification.....	7
7.2 Determination of moisture content.....	7
7.3 Determination of total ash content.....	7
7.4 Determination of acid-insoluble ash content.....	7
7.5 Determination of water-solution extractives content.....	7
7.6 TLC identification.....	7
7.7 Determination of polysaccharide content.....	7
7.8 Determination of marker compound content.....	7
7.9 Determination of heavy metals contents.....	7
7.10 Determination of pesticide residues contents.....	7
7.11 Determination of sulfur dioxide residues contents.....	8
8 Test report.....	8
9 Packaging, storage and transportation.....	8
10 Marking and labelling.....	8
Annex A (informative) Determination of water-soluble extractives.....	9
Annex B (informative) TLC identification.....	10
Annex C (informative) Determination of polysaccharide content.....	11
Annex D (informative) Determination of betaine content.....	13
Annex E (informative) Reference values of national and regional limits of moisture, total ash, water-soluble extractives, polysaccharides, betaine contents in <i>Lycium barbarum</i> and <i>Lycium chinense</i> fruit.....	16

Annex F (informative) Reference information and methods for differentiating <i>Lycium barbarum</i> L. and <i>Lycium chinense</i> Mill.	18
Bibliography	19

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 23193:2020

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html (standards.iteh.ai)

This document was prepared by Technical Committee ISO/TC 249, *Traditional Chinese medicine*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

Lycium barbarum and *Lycium chinense* fruit, commonly called Lycium fruit or *Lycii Fructus*, is the dried fruit of *Lycium barbarum* Linné or *Lycium chinense* Mill. (Fam. Solanaceae). Lycium fruit was firstly recorded in the book '*Divine Farmer's Classic of Materia Medica*', and it has a long history in China, Korea, Japan and other Southeast Asian nations, where it is used to nourish the liver and kidneys and replenish essence to improve vision. Clinically, owing to its medicinal properties, it plays an important role in the treatment of diseases such as immune suppression, cancer and diabetic retinopathy.

Additionally, *Lycium barbarum* and *Lycium chinense* fruit, with its sweet taste and warming property, is widely used in functional food and cosmetics. Lycium fruit and its finished products also have a very high reputation worldwide for their effectiveness, and account for a large market share in the international trade of Chinese herbal medicines.

Lycium barbarum and *Lycium chinense* fruit is widely cultivated in the northwest of China, Korea and Canada, among other places. However, the quality of Lycium fruit provided from different areas or by different cultivators is quite different. In addition, though *Lycium barbarum* and *Lycium chinense* fruit has been recorded in several pharmacopeia and standards, specifications and quality requirements in these standards vary. Thus, there is a clear and urgent need to develop an international standard for harmonizing the existing standards, as well as ensuring the safety and effectiveness of *Lycium barbarum* and *Lycium chinense* fruit.

As national implementation may differ, national standards bodies are invited to modify the values given in [5.3](#), [5.4](#), [5.5](#), [5.6](#), [5.8](#) and [Clause 9](#) in their national standards. Examples of national and regional values are given in [Annex E](#).

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 23193:2020

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>

Traditional Chinese medicine — *Lycium barbarum* and *Lycium chinense* fruit

1 Scope

This document specifies the minimum requirements and test methods for *Lycium barbarum* and *Lycium chinense* fruit, which is derived from the plant of *Lycium barbarum* L. or *Lycium chinense* Mill.

It is applicable to *Lycium barbarum* and *Lycium chinense* fruit that is sold and used as herbal raw materials in the international trade, including unprocessed and traditionally processed materials.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 1575, *Tea — Determination of total ash*

ISO 1577, *Tea — Determination of acid-insoluble ash*

ISO 18664, *Traditional Chinese Medicine — Determination of heavy metals in herbal medicines used in Traditional Chinese Medicine*

ISO 20409, *Traditional Chinese medicine — Panax notoginseng root and rhizome*

ISO 21371, *Traditional Chinese medicine — Labelling requirements of products intended for oral or topical use*

ISO 22258, *Traditional Chinese medicine — Determination of pesticide residues in natural products by gas chromatography*

ISO 22590, *Traditional Chinese medicine — Determination of sulfur dioxide in natural products by titration*

World Health Organization *Quality control methods for herbal materials*, 2011

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

— ISO Online browsing platform: available at <https://www.iso.org/obp>

— IEC Electropedia: available at <http://www.electropedia.org/>

3.1

***Lycium barbarum* fruit**

dried ripe fruit of *Lycium barbarum* L. (Fam. Solanaceae)

3.2

***Lycium chinense* fruit**

dried ripe fruit of *Lycium chinense* Mill. (Fam. Solanaceae)

3.3

batch

samples collected from the same particular place at the same time, of no more than 1 000 kg

3.4

final sample

samples for the test required in this document

Note 1 to entry: Final samples may be packed in different materials meeting conditions for specific tests (e.g. moisture or total ash).


4 Descriptions

The structure of *Lycium barbarum* L., *Lycium chinense* Mill. and the dried ripe fruit are shown in [Figure 1](#). Different features such as leaves, flowers and fruits in *Lycium barbarum* L. and *Lycium chinense* Mill., and methods for differentiating these two species, are given in [Annex F](#).

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 23193:2020

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>



a) Plant of *Lycium barbarum* L.

b) Plant of *Lycium Chinese* Mill.

STANDARD PREVIEW
 (standards.it/it,ai)

ISO 23193:2020
<https://standards.iteh.ai/catalog/standards/sist/904640-23193-3b-57a5-a455871198c/iso-23193-2020>



c) Dried ripe fruits (upper: *Lycium barbarum* fruit; lower: *Lycium chinense* fruit)

Key

- | | |
|------------------------------------|---------------------|
| 1 fruit spur | 5 pistil |
| 2 flower | 6 pistil stalk scar |
| 3 corolla expended to show stamens | 7 fruit stalk scar |
| 4 stamen | |

Figure 1 — Structure of *Lycium barbarum* and *Lycium chinense* fruit

5 Requirements

5.1 Morphological features

5.1.1 Appearance

The fruit is nearly fusiform or elliptical. Pericarp is soft and externally roughly wrinkled. Sarcocarp is pulpy, soft and tender.

5.1.2 Colour

The external surface is red or dark red with a pistil stalk scar (6) at the apex and a white fruit stalk scar (7) at the base, as in [Figure 1 c](#)).

5.1.3 Dimensions

The fruit is 6 mm to 20 mm in length measured from the base to the end of the fruit and 3 mm to 10 mm in diameter measured at the base of the fruit.

5.1.4 Fracture


20 to 50 seeds are present inside one fruit, kidney-shaped and nearly flat, ca. 2 mm; the external surface of the seeds is pale yellow or yellowish brown in the fruit of *Lycium barbarum* L., while the seeds in one fruit of *Lycium chinense* Mill. are numerous, kidney-shaped and nearly flat, 2,5 mm to 3 mm; the external surface colour of the seeds is yellow.

5.1.5 Odour

The odour is slight, and the taste is at first sweet and then slightly bitter for *Lycium chinense* fruit, and sweet and not bitter for *Lycium barbarum* fruit.

5.2 Microscopic characteristics

See [Figure 2](#). The powder is yellowish-orange or reddish-brown. Epidermal cells of outer pericarp (1) are polygonal or elongated-polygonal, about 60 µm in diameter, with straight or slightly wavy walls, covered with a thick cuticle, with distinct, more-or-less parallel striations. Parenchymatous cells of mesocarp (2) are thin-walled subpolygonal, containing reddish-orange or brownish-red spherical aleurone granules (3). Fragments of starchy endosperm cells (4) contain oil droplets. Stone cells of testa (5) are irregular polygonal, with distinct striations, thickened or slightly wavy walls.

**Key**

- 1 epidermal cells of outer pericarp
- 2 parenchymatous cells of mesocarp
- 3 aleurone granules
- 4 starchy endosperm cells
- 5 stone cells of testa

ISO 23193:2020

<https://standards.iteh.ai/catalog/standards/sist/77604640-23d9-482b-91a5-a45587725f8c/iso-23193-2020>

Figure 2 — Structure of powdered Lycium fruit**5.3 Moisture**

The mass fraction of moisture should not be more than 13,0 %.

5.4 Total ash

The mass fraction of total ash should not be more than 6,0 %.

5.5 Acid-insoluble ash

The mass fraction of total ash should not be more than 1,0 %.

5.6 Water-soluble extractives

The mass fraction of water-soluble extractives should not be less than 45,0 %.