



SLOVENSKI STANDARD
oSIST prEN ISO 18187:2023
01-september-2023

**Kakovost tal - Kontaktni preskus za trdne vzorce z dehidrogenazno aktivnostjo
Arthrobacter globiformis (ISO/DIS 18187:2023)**

Soil quality - Contact test for solid samples using the dehydrogenase activity of
Arthrobacter globiformis (ISO/DIS 18187:2023)

Bodenbeschaffenheit - Feststoffkontakttest unter Verwendung der
Dehydrogenaseaktivität von Arthrobacter globiformis (ISO/DIS 18187:2023)

Qualité du sol - Essai contact pour échantillons solides utilisant l'activité déshydrogénase
de Arthrobacter globiformis (ISO/DIS 18187:2023)

Ta slovenski standard je istoveten z: prEN ISO 18187

ICS:

13.080.30 Biološke lastnosti tal Biological properties of soils

oSIST prEN ISO 18187:2023 **en,fr,de**

DRAFT INTERNATIONAL STANDARD

ISO/DIS 18187

ISO/TC 190/SC 4

Secretariat: AFNOR

Voting begins on:
2023-06-29Voting terminates on:
2023-09-21

Soil quality — Contact test for solid samples using the dehydrogenase activity of *Arthrobacter globiformis*

*Qualité du sol — Essai contact pour échantillons solides utilisant l'activité déshydrogénase de *Arthrobacter globiformis**

ICS: 13.080.30

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN ISO 18187:2023](https://standards.iteh.ai/catalog/standards/sist/3f3814d8-f656-4302-a623-cabb300ad9bc/osist-pren-iso-18187-2023)

<https://standards.iteh.ai/catalog/standards/sist/3f3814d8-f656-4302-a623-cabb300ad9bc/osist-pren-iso-18187-2023>

This document is circulated as received from the committee secretariat.

THIS DOCUMENT IS A DRAFT CIRCULATED FOR COMMENT AND APPROVAL. IT IS THEREFORE SUBJECT TO CHANGE AND MAY NOT BE REFERRED TO AS AN INTERNATIONAL STANDARD UNTIL PUBLISHED AS SUCH.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

ISO/CEN PARALLEL PROCESSING



Reference number
ISO/DIS 18187:2023(E)

© ISO 2023

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN ISO 18187:2023](https://standards.iteh.ai/catalog/standards/sist/3f3814d8-f656-4302-a623-cabb300ad9bc/osist-pren-iso-18187-2023)

<https://standards.iteh.ai/catalog/standards/sist/3f3814d8-f656-4302-a623-cabb300ad9bc/osist-pren-iso-18187-2023>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2023

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

	Page
Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	4
5 Reagents and material	4
5.1 Test organisms.....	4
5.2 Control substrates.....	4
5.2.1 General.....	4
5.2.2 Control for soils.....	5
5.2.3 Control for waste material.....	5
5.3 Test substrates.....	5
5.4 Chemicals.....	6
6 Apparatus	8
7 Procedure	9
7.1 Preparation of dilutions.....	9
7.2 Reference substance and positive control preparation.....	9
7.3 Contact test procedure.....	10
7.3.1 General.....	10
7.3.2 Aeration.....	11
7.3.3 Deactivation.....	11
7.3.4 Preparation of the inoculum.....	11
7.3.5 Incubation and fluorescence measurement.....	11
7.4 Interferences.....	12
8 Calculation and expression of results	12
8.1 Calculation.....	12
8.1.1 Relative fluorescence.....	12
8.1.2 Determining the percentage of inhibition.....	12
8.2 Expression of results.....	13
9 Validity of the test	13
10 Statistical analysis	14
11 Test report	14
Annex A (informative) Results on the ring test	16
Annex B (informative) Preparation of test organisms	23
Annex C (informative) Testing chemical substances	25
Bibliography	27

ISO/DIS 18187:2023(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO document should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

ISO draws attention to the possibility that the implementation of this document may involve the use of (a) patent(s). ISO takes no position concerning the evidence, validity or applicability of any claimed patent rights in respect thereof. As of the date of publication of this document, ISO [had/had not] received notice of (a) patent(s) which may be required to implement this document. However, implementers are cautioned that this may not represent the latest information, which may be obtained from the patent database available at www.iso.org/patents. ISO shall not be held responsible for identifying any or all such patent rights.

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

This second/third/... edition cancels and replaces the first/second/... edition (ISO #####:#####), which has been technically revised.

The main changes are as follows:

— xxx xxxxxxxx xxx xxxxx

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

This International Standard describes the miniaturized solid contact assay with *Arthrobacter globiformis* that allows the preliminary assessment of solid material (i.e. soil and waste materials) within 6 h. The principle of the assay relies on dehydrogenase activity inhibition of an added test organism, caused by bioavailable toxic substances in soil and waste samples. This is an ecologically relevant assay as far as it uses a ubiquitous soil bacterial species with high affinity to surfaces [15] [6] which dehydrogenases are involved in different biological mechanisms withstanding bacteria integrity (e.g. respiratory chains). Moreover, it has been noticed that this parameter (dehydrogenase activity inhibition) is quite sensitive to different toxic substances [10] [12] [14] [15] [19].

Overall, this assay is non-labour-intensive, rapid, cost-effective and sensitive, providing results that improve the physical and chemical assessment of natural samples while allowing a quick indication of their biological effects.

The miniaturized solid contact assay is based on the solid contact assay established by Reference [7].

This International Standard is also based on Reference [23].

The results of an interlaboratory trial towards the evaluation of test variability to assess different waste and soil samples, as well as chemicals, are presented in [Annex A](#) and in Reference [33].

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN ISO 18187:2023](#)

<https://standards.iteh.ai/catalog/standards/sist/3f3814d8-f656-4302-a623-cabb300ad9bc/osist-pren-iso-18187-2023>

Soil quality — Contact test for solid samples using the dehydrogenase activity of *Arthrobacter globiformis*

1 Scope

This International Standard specifies a rapid method for assessing solid samples in an aerobic suspension, by determining the inhibition of dehydrogenase activity of *Arthrobacter globiformis* using the redox dye resazurin.

It is applicable for assessing the effect of water-soluble and solid matter bounded non-volatile contaminants of natural samples, such as soils and waste materials. Although not the main purpose, the contact test can additionally be used for testing the effect of chemicals, as described in the [Annex C](#). The test yields a result within 6 h and can therefore be used for screening potentially contaminated material.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 5667-15, *Water quality — Sampling — Part 15: Guidance on the preservation and handling of sludge and sediment samples*

ISO 10381-6, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*

CEN/TR 15310-1, *Characterization of waste — Sampling of waste materials — Part 1: Guidance on selection and application of criteria for sampling under various conditions*

EN 14735, *Characterization of waste — Preparation of waste samples for ecotoxicity tests*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.1

contact time

exposure period of the bacteria to a suspension of solid matter

3.2

negative control

sample of a *control substrate* (3.6) with a mixture of known solutions (distilled water, medium B or *inoculum* (3.12))

Note 1 to entry: It is used to standardize the analysis.

ISO/DIS 18187:2023(E)

3.3

positive control

sample of a *control substrate* (3.6) with a mixture of known solutions [distilled water, medium B or *inoculum* (3.12)] and a reference substance

Note 1 to entry: It is used to check the sensitivity of the test organism.

3.4

blank A

blank, which sets the own fluorescence of the substrate after being deactivated

Note 1 to entry: Blank A is not added with bacteria.

3.5

blank B

blank, which sets the natural fluorescence of the substrate without being deactivated

Note 1 to entry: Blank B is not added with bacteria.

3.6

control substrate

reference or standard substrate used as a control and as a dilution substrate for preparing dilution/concentration series with *test substrates* (3.7) or a reference substance

EXAMPLE Quartz sand or LUFA standard soil type 2.2.

3.7

test substrate

natural or artificial substrate that is naturally contaminated or spiked with a test chemical

Note 1 to entry: The test substrate is the *test material* (3.8) after being prepared for testing (e.g. sieved) and/or diluted with a *control substrate* (3.6).

3.8

test material

original sample of soil or waste material without any changes (e.g. sieving)

3.9

dehydrogenase activity

activity of hydrogen-abstracting enzymes which are involved in many energy, redox reactions, and biosynthetic metabolic processes (e.g. the respiratory chain), which require cell integrity to be produced

Note 1 to entry: These enzymes can reduce resazurin into resorufin in the extracellular environment^[6].

Note 2 to entry: See Reference [21].

3.10

effect concentration for x % effect**EC_x**

concentration (mass fraction) of a test substance or sample that causes x % of an effect on a given endpoint within a given exposure period when compared with a control

EXAMPLE An EC₅₀ is a concentration estimated to cause an effect on a test end point in 50 % of an exposed population over a defined exposure period.

Note 1 to entry: The EC_x is expressed as a percentage of soil or waste tested per dry mass of soil mixture. When chemicals are tested, the EC_x is expressed as mass of the test substance per dry mass of soil, in milligrams per kilogram.

3.11**freeze-dried bacteria**

bacterial culture preserved through the water removing of a frozen cell suspension by sublimation under reduced vacuum pressure

Note 1 to entry: The preserved cultures can be stored at (-20 ± 2) °C. The bacteria are active after being reconstituted with sterilized distilled water [20 min to 30 min at (6 ± 2) °C] and ready to be used in the test, see [7.3.4 b](#)).

3.12**inoculum**

suspension of bacteria used to inoculate a nutrient solution

3.13**medium**

aqueous nutritive solution required for bacterial growth

3.14**optical density of bacterial inoculum**

measurement of the attenuation of a light beam passing through a bacterial suspension at 600 nm (used to determine the cell count indirectly)

Note 1 to entry: In a bacterial test, the absorbance is usually measured as FAU (formazine attenuation units) at 600 nm (see Reference [\[3\]](#)).

3.15**test start**

moment when the substrates, reagents and the bacterial *inoculum* ([3.12](#)) are prepared immediately before the incubation and reaction period

Note 1 to entry: Here is when preparing the test and *control substrates* ([3.6](#)) for incubation (i.e. [Table 1](#), day 0).

3.16**reaction time**

time it takes for the enzyme to react (from the addition of the resazurin solution until the end of the kinetics period)

3.17**slope**

quotient of the *relative fluorescence* ([3.18](#)) variation along the *reaction time* ([3.16](#)) between 15 min and 45 min

Note 1 to entry: The slope (expressed as min^{-1}) results from fitting a linear regression model to the fluorescence readings over time.

3.18**relative fluorescence**

fluorescence measured for each treatment (control and test) after subtracting the fluorescence of the respective *blank A* ([3.4](#))

3.19**stock culture**

bacterial culture obtained from a pure strain culture acquired from a certified laboratory

Note 1 to entry: This stock culture provides an *inoculum* ([3.12](#)) for the pre-culture in the test procedure.

ISO/DIS 18187:2023(E)

3.20

lowest ineffective dilution

LID-value

lowest value of the dilution factor (LID) for which the test does not give an ecotoxicological relevant reduction

Note 1 to entry: The LID is expressed as the reciprocal value of dilution.

EXAMPLE An often used dilution series is 1/2/4/8/16 [= 100 %/50 %/25 %/12,5 %/6,25 % *test substrate* (3.7) to *control substrate* (3.6)]. A LID 8 corresponds to a dilution of soil or waste of 1:8.

4 Principle

The bacteria *Arthrobacter globiformis* is added to the solid material and incubated at $(30 \pm 1) ^\circ\text{C}$ for 2 h. After this contact time, the non-toxic redox dye resazurin is added. Due to the dehydrogenase activity, resazurin is transformed into resorufin, in the extracellular environment.^[6] Resorufin can be detected fluorometrically (excitation at 535 nm, emission at 590 nm) in the presence of solid matter. The increase of resorufin is determined by measuring the fluorescence every 15 min for a period of 1 h. In order to determine the inhibition of the dehydrogenase activity, the rate of resorufin increase in the sample is compared with the rate of resorufin increase in the control. In the presence of toxic substances, an inhibition of dehydrogenase activity is expected. This is reflected by the reduction of resorufin production and subsequent lowering of fluorescence emission.

5 Reagents and material

5.1 Test organisms

The test organism is *Arthrobacter globiformis* (Conn 1982) Conn and Dimmick 1947 (strain number ATCC 8010TM *Arthrobacter globiformis* (Conn) Conn and Dimmick), which is common in soils. *Arthrobacter* species belong to the *Micrococaceae* family. They are mostly obligate aerobic organisms, although some species can exhibit anaerobic metabolism under limiting oxygen conditions.^[9] *Arthrobacter* spp. are chemoheterotrophic, and present pleomorphic characteristics, since they show a rod-to-coccus morphology change as they enter in the stationary phase. Although *Arthrobacter* is gram-positive, it can stain gram-negative during the log-phase. Variations in the cell wall thickness along the bacteria growth can lead to gram variability by differential staining of the granules.^[22] However, this characteristic does not induce a differential sensitivity between assays, as far as an inoculum in exponential growth phase is used during the reaction time. *Arthrobacter globiformis* is classified in the risk group I — non-pathogenic organism.

The bacteria strain can be achieved from commercially available freeze-dried or liquid-dried reagents, or from culture collections, e.g. Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH (strain number 20124), or ARS Culture collection NCAUR.¹⁾ The bacterial suspensions used for toxicity measurements shall be freshly prepared from stock cultures or directly used from a ready-to-use freeze-dried batch. The stock culturing and freeze-drying process of the bacteria is described in [Annex B](#).

5.2 Control substrates

5.2.1 General

The control substrates selected from the options presented below are to prepare both the negative (addition of distilled water, see [5.2.2](#), [5.2.3](#)) and positive (addition of the reference substance, see [7.2](#))

1) Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ) GmbH, Mascheroder Weg 10, D-38124 Braunschweig, Germany; or ARS (Agricultural Research Service) Culture collection (also known as NRRL) belonging to the National Center for Agricultural Utilization Research (NCAUR), 1815 N, University Street, Peoria, Illinois 61604, USA are examples of firms that sell this bacteria. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these firms.