
Kemična razkužila in antiseptiki - Kvantitativni suspenzijski preskus za vrednotenje fungicidnega delovanja ali delovanja kemičnih razkužil in antiseptikov na kvasovke v veterini - Preskusna metoda in zahteve (faza 2, stopnja 1)

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of fungicidal or yeasticidal activity of chemical disinfectants and antiseptics used in the veterinary area - Test method and requirements (phase 2, step 1)

Chemische Desinfektionsmittel und Antiseptika - Quantitativer Suspensionsversuch zur Bestimmung der fungiziden oder levuroziden Wirkung chemischer Desinfektionsmittel und Antiseptika für den Veterinärbereich - Prüfverfahren und Anforderungen (Phase 2, Stufe 1)

Antiseptiques et désinfectants chimiques - Essai quantitatif de suspension pour l'évaluation de l'activité fongicide ou levuricide des antiseptiques et des désinfectants chimiques utilisés dans le domaine vétérinaire - Méthode d'essai et prescriptions (ph

Ta slovenski standard je istoveten z: EN 1657:2024

ICS:

11.080.20	Dezinfektanti in antiseptiki	Disinfectants and antiseptics
11.220	Veterinarstvo	Veterinary medicine

SIST EN 1657:2025**en,fr,de**

EUROPEAN STANDARD

EN 1657

NORME EUROPÉENNE

EUROPÄISCHE NORM

December 2024

ICS 71.100.35

Supersedes EN 1657:2016

English Version

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of fungicidal or yeasticidal activity of chemical disinfectants and antiseptics used in the veterinary area - Test method and requirements (phase 2, step 1)

Antiseptiques et désinfectants chimiques - Essai quantitatif de suspension pour l'évaluation de l'activité fongicide ou levuricide des antiseptiques et des désinfectants chimiques utilisés dans le domaine vétérinaire - Méthode d'essai et exigences (phase 2, étape 1)

Chemische Desinfektionsmittel und Antiseptika - Quantitativer Suspensionsversuch zur Bestimmung der fungiziden oder levuroziden Wirkung chemischer Desinfektionsmittel und Antiseptika für den Veterinärbereich - Prüfverfahren und Anforderungen (Phase 2, Stufe 1)

This European Standard was approved by CEN on 18 November 2024.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Türkiye and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

Contents

Page

European foreword.....	4
Introduction	5
1 Scope	6
2 Normative references	6
3 Terms, definitions and abbreviated terms	6
3.1 Terms and definitions	6
3.2 Symbols and abbreviated terms	7
4 Requirements	7
5 Test method	8
5.1 Principle	8
5.2 Materials and reagents.....	9
5.2.1 Test organisms	9
5.2.2 Culture media and reagents	9
5.3 Apparatus and glassware	12
5.3.1 General.....	12
5.3.2 Usual microbiological laboratory equipment and, in particular, the following.....	12
5.4 Preparation of test organism suspensions and product test solutions.....	13
5.4.1 Test organism suspensions (test and validation suspension).....	13
5.4.2 Product test solutions.....	18
5.5 Procedure for assessing the fungicidal or yeasticidal activity of the product.....	18
5.5.1 General.....	18
5.5.2 Dilution-neutralization method	19
5.5.3 Membrane filtration method	22
5.6 Experimental data and calculation.....	24
5.6.1 Explanation of terms and abbreviations	24
5.6.2 Calculation.....	24
5.7 Verification of methodology.....	27
5.7.1 General.....	27
5.7.2 Control of weighted mean counts.....	27
5.7.3 Basic limits	28
5.7.4 Additional limits for <i>Aspergillus brasiliensis</i>	28
5.8 Expression of results and precision	28
5.8.1 Reduction.....	28
5.8.2 Control of active and non-active product test solution (5.4.2).....	28
5.8.3 Limiting test organism and fungicidal/yeasticidal concentration.....	28
5.8.4 Precision, replicates.....	29
5.9 Interpretation of results - conclusion	29
5.9.1 General.....	29
5.9.2 Fungicidal activity for general purposes.....	29
5.9.3 Fungicidal activity for specific purposes	29
5.9.4 Yeasticidal activity for general purposes.....	29
5.9.5 Yeasticidal activity for specific purposes	29
5.9.6 Yeasticidal activity for teat disinfectants	30
5.10 Test report.....	30

Annex A (informative) Referenced strains in national collections	32
Annex B (informative) Neutralizers and rinsing liquids	33
Annex C (informative) Graphical representation of test procedures	35
C.1 Dilution-neutralization method	35
C.2 Membrane filtration method	37
Annex D (informative) Example of a typical test report	40
Annex E (informative) Precision of the test result	45
Bibliography	48

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[SIST EN 1657:2025](https://standards.iteh.ai/catalog/standards/sist/525b6dca-1ccf-48ba-ae3b-d264bca3a931/sist-en-1657-2025)

<https://standards.iteh.ai/catalog/standards/sist/525b6dca-1ccf-48ba-ae3b-d264bca3a931/sist-en-1657-2025>

EN 1657:2024 (E)**European foreword**

This document (EN 1657:2024) has been prepared by Technical Committee CEN/TC 216 “Chemical disinfectants and antiseptics”, the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by June 2025, and conflicting national standards shall be withdrawn at the latest by June 2025.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 1657:2016.

This document was revised to correct obvious errors and ambiguities, to harmonize the structure and wording with other tests of CEN/TC 216 (existing or in preparation), and to improve the readability of the standard and thereby make it more understandable.

The obligatory conditions have been removed and the preparation of the milk interfering substance has been amended.

The data obtained using the 2016 version of EN 1657 are still valid.

Any feedback and questions on this document should be directed to the users' national standards body. A complete listing of these bodies can be found on the CEN website.

According to the CEN-CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Türkiye and the United Kingdom.

Introduction

This document specifies a suspension test method for establishing whether a chemical disinfectant or antiseptic has a fungicidal or yeasticidal activity in the areas described in the scope.

This laboratory test takes into account practical conditions of application of the product, including contact time, temperature, test organisms and interfering substances, i.e. conditions which may influence its action in practical situations.

The conditions are intended to cover general purposes and to allow reference between laboratories and product types. Each utilization concentration of the chemical disinfectant or antiseptic found by this test corresponds to defined experimental conditions. However, for some applications the recommendations of use of a product may differ and therefore additional test conditions need to be used.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[SIST EN 1657:2025](https://standards.iteh.ai/catalog/standards/sist/525b6dca-1ccf-48ba-ae3b-d264bca3a931/sist-en-1657-2025)

<https://standards.iteh.ai/catalog/standards/sist/525b6dca-1ccf-48ba-ae3b-d264bca3a931/sist-en-1657-2025>

EN 1657:2024 (E)**1 Scope**

This document specifies a test method and the minimum requirements for fungicidal or yeasticidal activity of chemical disinfectant and antiseptic products that form a homogeneous, physically stable preparation when diluted with hard water or — in the case of ready-to-use-products — with water. Products can only be tested at a concentration of 80 % or less, as some dilution is always produced by adding the test organisms and interfering substance.

This document applies to products that are used in the veterinary area – i.e. in the breeding, husbandry, production, veterinary care facilities, transport and disposal of all animals except when in the food chain following death and entry into processing industry. This document also applies to products used for teat disinfection.

EN 14885 specifies in detail the relationship of the various tests to one another and to “use recommendations”.

NOTE 1 The method described is intended to determine the activity of commercial formulations or active substances under the conditions in which they are used.

NOTE 2 This method corresponds to a phase 2 step 1 test.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 12353, *Chemical disinfectants and antiseptics — Preservation of test organisms used for the determination of bactericidal (including Legionella), mycobactericidal, sporicidal, fungicidal and virucidal (including bacteriophages) activity*

EN 14885, *Chemical disinfectants and antiseptics — Application of European Standards for chemical disinfectants and antiseptics*

ISO 4793, *Laboratory sintered (fritted) filters — Porosity grading, classification and designation*

3 Terms, definitions and abbreviated terms**3.1 Terms and definitions**

For the purposes of this document, the terms and definitions given in EN 14885 apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

- ISO Online Browsing Platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.2 Symbols and abbreviated terms

For the purposes of this document, the following symbols and abbreviated terms apply.

c	is the sum of V_C values taken into account
cfu	Colony forming units
n	is the number of V_C values taken into account
N	Test Suspension
N_v	Validation suspension
N_a	Test
A	Control A (Experimental conditions control)
B	Control B (Neutralization control)
C	Control C (Dilution neutralization validation)
N_0	Test at time 0
N_{v0}	Validation suspension at time 0
R	Reduction
V_c	is the number of cfu counted per sample of 1,0ml

4 Requirements

The product shall demonstrate at least a 4 decimal log (lg) reduction when diluted with hard water (5.2.2.7) or – in the case of ready-to-use products – with water (5.2.2.2) and tested in accordance with Table 1 and Clause 5 under simulated low-level soiling (3,0 g/l bovine albumin) or high level soiling (10,0 g/l yeast extract and 10,0 g/l bovine albumin) or 10,0 g/l milk for teat disinfectants or 3,0 g/l bovine albumin for pre-milking teat disinfectants in additional test conditions.

Table 1 — Test conditions

Test conditions	Fungicidal activity	Yeasticidal activity	Yeasticidal activity for teat disinfectants
Test organisms	<i>Aspergillus brasiliensis</i> <i>Candida albicans</i>	<i>Candida albicans</i>	<i>Candida albicans</i>
additional	any relevant test organism	any relevant test organism	any relevant test organism
Test temperature	At intervals of 5°C		
Minimum	5°C ± 1°C	5°C ± 1°C	20°C ± 1°C
Maximum	40°C ± 1°C	40°C ± 1°C	30°C ± 1°C
Contact time	At intervals of 30 s from 30 s to 5 min and at intervals of 5 min from 5 min to 120 min		
Minimum			
Maximum	1 min ± 5 s	1 min ± 5 s	1 min ± 5 s for post-milking teat disinfectants 30 s ± 5 s for pre-milking teat disinfectants
	120 min ± 10 s	120 min ± 10 s	5 min ± 10 s for post-milking teat disinfectants 3 min ± 10 s for pre-milking teat disinfectants
Interfering substance	Document Preview		
low level soiling	3,0 g/l bovine albumin	3,0 g/l bovine albumin	Post milking: 10,0 g/l of reconstituted milk
high level soiling	10 g/l yeast extract plus 10 g/l bovine albumin	10 g/l yeast extract plus 10 g/l bovine albumin	Pre-milking: 3,0 g/l bovine albumin
additional	any relevant substance	any relevant substance	any relevant substance
NOTE For the additional conditions, the concentration defined as a result can be lower than the one obtained under the standard test conditions.			
The recommended test conditions for the use of the product are within the responsibility of the manufacturer.			

Any additional specific fungicidal or yeasticidal activity shall be determined in accordance with 5.2.1 and 5.5.1.1 in order to take into account intended specific use conditions.

5 Test method

5.1 Principle

5.1.1 A sample of the product as delivered and/or diluted with hard water (or water for ready-to-use products) is added to a test suspension of fungi (yeast cells or mould spores) in a solution of an interfering substance. The mixture is maintained at temperature θ for the test contact time t in accordance with Table 1. At the end of this contact time, an aliquot is taken, and the fungicidal/yeasticidal and/or the fungistatic/yeastistatic activity in this portion is immediately neutralized or suppressed by a validated

method. The method of choice is dilution-neutralization. If a suitable neutralizer cannot be found, membrane filtration is used. The number of surviving fungi in each sample are determined and the reduction is calculated.

5.2 Materials and reagents

5.2.1 Test organisms

The fungicidal activity shall be evaluated using the following strains as test organisms:

- *Candida albicans*;
- *Aspergillus brasiliensis*.

(formerly *Aspergillus niger*)

The yeasticidal activity shall be evaluated using only *Candida albicans*.

NOTE See Annex A for strain references in some culture collections.

The required incubation temperature for these test organisms is $30\text{ °C} \pm 1\text{ °C}$ (see 5.3.2.3). The same temperature shall be used for all incubations performed during a test and its control and validation.

If additional test organisms are used, they shall be kept and used under optimum growth conditions (temperature, time, atmosphere, media) and noted in the test report. If the additional test organisms selected do not correspond to the specified strains, their suitability for supplying the required inocula shall be verified. If these additional test organisms are not classified at a reference centre, their identification characteristics shall be stated. In addition, they shall be held by the testing laboratory or national culture collection under a reference for five years.

5.2.2 Culture media and reagents

5.2.2.1 General

SIST EN 1657:2025

<https://standards.iteh.ai/catalog/standards/sist/525b6dca-1ccf-48ba-ae3b-d264bca3a931/sist-en-1657-2025>

All weights of chemical substances given in this Standard refer to the anhydrous salts. Hydrated forms may be used as an alternative, but the weights required shall be adjusted to allow for consequent molecular weight differences.

The reagents shall be of analytical grade and/or appropriate for microbiological purposes. They shall be free from substances that are toxic or inhibitory to the test organisms.

If additional strains do not grow on the media (5.2.2.3) or cannot be used with diluent (5.2.2.4) alternative media shall be used and shall be reported as well as alternative incubation conditions.

To improve reproducibility, it is recommended that commercially available dehydrated material (if appropriate) is used for the preparation of culture media. The manufacturer's instructions relating to the preparation of these products should be rigorously followed.

Ready to use media may be used if it complies with the required specification.

For each culture medium and reagent, a time limitation for use should be fixed.

All specified pH values are measured at $20\text{ °C} \pm 1\text{ °C}$.

5.2.2.2 Water

The water shall be freshly glass-distilled water and not demineralized water. If distilled water of adequate quality is not available, water for injections (see bibliographic reference [1]) can be used.

EN 1657:2024 (E)

Sterilize in the autoclave [5.3.2.1 a)]. Sterilization is not necessary if the water is used e.g. for preparation of culture media and subsequently sterilized.

NOTE 1 See 5.2.2.7 for the procedure to prepare hard water.

5.2.2.3 Malt extract agar (MEA)

Malt extract agar, consisting of:

Malt extract ^a	30,0 g
Agar	15,0 g
Water (5.2.2.2)	to 1 000,0 ml

^a The malt extract should be of food grade (Cristomalt poudre from Difal is recommended) or equivalent that is not highly purified and not only based on maltose (Malt extract from OXOID is recommended)¹.

Sterilize in the autoclave [5.3.2.1 a)]. After sterilization, the pH of the medium shall be equivalent to $5,6 \pm 0,2$ when measured at $20 \text{ }^\circ\text{C} \pm 1 \text{ }^\circ\text{C}$.

In case of encountering problems with neutralization (5.5.1.2 and 5.5.1.3), it may be necessary to add neutralizer to the MEA. Annex B gives guidance on the neutralizers that may be used.

5.2.2.4 Diluent

Tryptone sodium chloride solution, consisting of:

Tryptone, pancreatic digest of casein	1,0 g
Sodium chloride (NaCl)	8,5 g
Water (5.2.2.2)	to 1 000,0 ml

Sterilize in the autoclave [5.3.2.1 a)]. After sterilization, the pH of the diluent shall be equivalent to $7,0 \pm 0,2$ when measured at $20 \text{ }^\circ\text{C} \pm 1 \text{ }^\circ\text{C}$.

5.2.2.5 Neutralizer

The neutralizer shall be validated for the product being tested in accordance with 5.5.1.2, 5.5.1.3 and 5.5.2. It shall be sterile.

NOTE Information on neutralizers that have been found to be suitable for some categories of products is given in Annex B.

5.2.2.6 Rinsing liquid (for membrane filtration)

The rinsing liquid shall be validated for the product being tested in accordance with 5.5.1.2, 5.5.1.3 and 5.5.3. It shall be sterile, compatible with the filter membrane and capable of filtration through the filter membrane under the test conditions described in 5.5.3.

NOTE Information on rinsing liquids that have been found to be suitable for some categories of products is given in Annex B.

5.2.2.7 Hard water for dilution of products

For the preparation of 1 l of hard water, the procedure is as follows:

¹ This information is given for the convenience of users of this European Standard and does not constitute an endorsement by CEN of the product named. Equivalent products may be used if they can be shown to lead to the same results.

- prepare solution A: dissolve 19,84 g magnesium chloride (MgCl_2) and 46,24 g calcium chloride (CaCl_2) in water (5.2.2.2) and dilute to 1 000 ml. Sterilize by membrane filtration (5.3.2.7) or in the autoclave [5.3.2.1 a)].

Autoclaving – if used – may cause a loss of liquid. In this case make up to 1 000 ml with water (5.2.2.2) under aseptic conditions. Store the solution in the refrigerator (5.3.2.8) for no longer than one month;

- prepare solution B: dissolve 35,02 g sodium bicarbonate (NaHCO_3) in water (5.2.2.2) and dilute to 1 000 ml.
- Sterilize by membrane filtration (5.3.2.7). Store the solution in the refrigerator (5.3.2.8) for no longer than one week;
- place 600 ml to 700 ml of water (5.2.2.2) in a 1 000 ml volumetric flask (5.3.2.12) and add 6,0 ml (5.3.2.9) of solution A, then 8,0 ml of solution B. Mix and dilute to 1 000 ml with water (5.2.2.2). The pH of the hard water shall be $7,0 \pm 0,2$, when measured at $20 \text{ °C} \pm 1 \text{ °C}$ (5.3.2.4). If necessary, adjust the pH by using a solution of approximately 40 g/l (about 1 mol/l) of sodium hydroxide (NaOH) or approximately 36,5 g/l (about 1 mol/l) of hydrochloric acid (HCl).

The hard water shall be freshly prepared under aseptic conditions and used within 12 h.

NOTE When preparing the product test solutions (5.4.2), the addition of the product to the hard water produces a different final water hardness in each test tube. In any case the final hardness is lower than 375 mg/l of calcium carbonate (CaCO_3) in the test tube.

5.2.2.8 Interfering substances

5.2.2.8.1 General

The interfering substance shall be chosen according to the conditions of use laid down for the product.

The interfering substance shall be sterile and prepared at 10 times its final concentration in the test.

For any additional interfering substance the ionic composition (e.g. pH, calcium and/or magnesium hardness) and chemical composition (e.g. mineral substances, protein, carbohydrates, lipids and detergents) shall be defined.

NOTE The term “interfering substance” is used even if it contains more than one substance.

5.2.2.8.2 Low-level and pre-milking soiling (bovine albumin solution)

Dissolve 3,0 g of bovine albumin fraction V (suitable for microbiological purposes) in 90 ml of water (5.2.2.2) in a 100ml volumetric flask (5.3.2.12). Make up to the mark with water.

Sterilize by membrane filtration (5.3.2.7), keep in the refrigerator (5.3.2.8) and use within one month.

The final concentration of bovine albumin in the test procedure (5.5) is 3,0 g/l.

5.2.2.8.3 High-level soiling (mixture of bovine albumin solution with yeast extract)

Dissolve 50,0 g yeast extract powder in 150 ml of water (5.2.2.2) in a 250 ml volumetric flask (5.3.2.12) and allow foam to collapse. Make up to the mark with water (5.2.2.2). Transfer to a clean dry bottle and sterilize in an autoclave [5.3.2.1 a)]. Allow to cool to $20 \text{ °C} \pm 1 \text{ °C}$.

Pipette 25 ml of this solution into a 50 ml volumetric flask (5.3.2.12) and add 10 ml of water (5.2.2.2). Dissolve 5,0 g of bovine albumin fraction V (suitable for microbiological purposes) in the solution with

EN 1657:2024 (E)

shaking and allow foam to collapse. Make up to the mark with water (5.2.2.2), sterilize by membrane filtration (5.3.2.7), keep in the refrigerator (5.3.2.8) and use within one month.

The final concentration in the test procedure (5.5) is 10,0 g/l yeast extract and 10,0 g/l bovine albumin.

5.2.2.8.4 Milk for teat disinfectants

Skimmed milk, guaranteed free of antibiotics and additives and reconstituted at a rate of 100 g powder per litre of water (5.2.2.2), shall be prepared as follows:

- prepare a solution of 10,0 % (v/v) in water (5.2.2.2) by adding 10 parts of reconstituted milk to 90 parts of water. Heat for 30 min at $(105 \pm 3) ^\circ\text{C}$ [or 5 min at $(121 \pm 3) ^\circ\text{C}$].

The final concentration of reconstituted milk in the test procedure (5.5) is 1,0 % (v/v) of reconstituted milk.

5.3 Apparatus and glassware**5.3.1 General**

Sterilize all glassware and parts of the apparatus that will come into contact with the culture media and reagents or the sample, except those which are supplied sterile, by one of the following methods:

- a) by moist heat, in the autoclave [5.3.2.1 a)];
- b) by dry heat, in the hot air oven [5.3.2.1 b)].

5.3.2 Usual microbiological laboratory equipment² and, in particular, the following**5.3.2.1 Apparatus for sterilization**

- a) for moist heat sterilization, an autoclave capable of being maintained at $(121 \text{ }_0^{+3}) ^\circ\text{C}$ for a minimum holding time of 15 min;
- b) for dry heat sterilization, a hot air oven capable of being maintained at $(180 \text{ }_0^{+5}) ^\circ\text{C}$ for a minimum holding time of 30 min, at $(170 \text{ }_0^{+5}) ^\circ\text{C}$ for a minimum holding time of 1 h or at $(160 \text{ }_0^{+5}) ^\circ\text{C}$ for a minimum holding time of 2 h.

5.3.2.2 Water baths, capable of being controlled at the specified temperatures and at $45 ^\circ\text{C} \pm 1 ^\circ\text{C}$ (to maintain melted MEA in case of pour plate technique).

5.3.2.3 Incubator, capable of being controlled at the specified temperatures $\pm 1 ^\circ\text{C}$.

5.3.2.4 pH-meter, having an inaccuracy of calibration of no more than $\pm 0,1$ pH units at $20 ^\circ\text{C} \pm 1 ^\circ\text{C}$. A puncture electrode or a flat membrane electrode should be used for measuring the pH of the agar media (5.2.2.3).

5.3.2.5 Stopwatch

² Disposable sterile equipment is an acceptable alternative to reusable glassware.