



Designation: D5562 – 95 (Reapproved 2011)

Standard Test Method for Determination of the Acidity as Free Fatty Acids/Acid Number in the Presence of Ammonium or Triethanolamine Soaps¹

This standard is issued under the fixed designation D5562; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This test method covers determination of the acidity as free fatty acids in the presence of ammonium or triethanolamine soaps, or both, existing in a sample of sulfonated or sulfated oil, or both, by calculation from the total alkalinity and free-and-alkali combined fatty acids.

1.2 The values stated in SI units are to be regarded as standard.

1.3 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.*

2. Referenced Documents

2.1 *ASTM Standards:*²

D5350 Test Method for Determination of Organically Combined Sulfuric Anhydride by Titration, Test Method A

3. Significance and Use

3.1 This test method is intended for use in the determination of the acid number of sulfonated and sulfated oils for the purpose of quality assurance.

4. Apparatus

4.1 The apparatus required consists of a glass flask provided with a glass stopper and an air condenser. The connection between the flask and the condenser shall be a ground joint. Perforated glass beads shall be used to prevent bumping.

4.1.1 *Flask*—An Erlenmeyer flask (Fig. 1) made of a borosilicate glass, having a capacity of approximately 300 mL and provided with a glass stopper.

4.1.2 *Condenser*, consisting of a glass tube, 915 mm (36 in.) in length and 8 mm ($\frac{5}{16}$ in.) in outside diameter. The lower end of the tube shall be flared and ground to fit the mouth of the Erlenmeyer flask.

4.1.3 *Glass Beads*—Perforated glass beads, made of chemically resistant glass, approximately 4 mm ($\frac{5}{32}$ in.) in diameter. Before using, the glass beads shall be boiled thoroughly in several portions of water or until the wash water reacts neutral to a methyl orange indicator.

5. Reagents

5.1 *Ethyl Ether*.

5.2 *Methyl Orange Indicator Solution (1 g/L)*—Dissolve 0.1 g of methyl orange in 100 mL of water.

5.3 *Sodium Chloride (NaCl)*, solid.

5.4 *Sodium Hydroxide, Standard Solution (1 N)*—Accurately prepare and standardize a 1 N sodium hydroxide (NaOH) solution. Express the strength or concentration of the solution as mg of KOH per mL; 1 mL of 1 N NaOH solution is equivalent to 56.1 mg of KOH.

5.5 *Sodium Hydroxide, Standard Solution (0.5 N)*—Accurately prepare and standardize a 0.5 N NaOH solution. Express the strength of the solution as mg of KOH per mL; 1 mL of 0.5 N NaOH solution is equivalent to 28.05 mg of KOH.

5.6 *Sulfuric Acid, Standard (0.5 N)*—Accurately prepare and standardize a 0.5 N sulfuric acid (H_2SO_4) solution. Express the strength of the solution as mg of KOH per mL; 1 mL of 0.5 N H_2SO_4 is equivalent to 28.05 mg of KOH.

5.7 *Sulfuric Acid (1 + 19)*—Carefully mix 1 volume of concentrated sulfuric acid (H_2SO_4 , sp gr 1.84) into 19 volumes of water while stirring.

6. Procedure

6.1 The procedure consists of two determinations: (1) total alkalinity and (2) free-and-alkali-combined fatty acids.

¹ This test method is under the jurisdiction of ASTM Committee D31 on Leather and is the direct responsibility of Subcommittee D31.08 on Fats and Oils. This test method was developed in cooperation with the American Leather Chemists Assn. (Method H 52-1956).

Current edition approved Jan. 1, 2011. Published March 2011. Originally approved in 1994. Last previous edition approved in 2006 as D5562 – 95(2006). DOI: 10.1520/D5562-95R11.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.