
International Standard



2962

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION • МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ • ORGANISATION INTERNATIONALE DE NORMALISATION

Cheese and processed cheese products — Determination of total phosphorus content — Molecular absorption spectrometric method

Fromages et fromages fondus — Détermination de la teneur en phosphore total — Méthode par spectrométrie d'absorption moléculaire

iTeh STANDARD PREVIEW

Second edition — 1984-06-01 (standards.iteh.ai)

[ISO 2962:1984](#)

<https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984>

UDC 637.3 : 543.42 : 546.18

Ref. No. ISO 2962-1984 (E)

Descriptors : agricultural products, dairy products, cheeses, processed cheese, chemical analysis, determination of content, phosphorus, spectrophotometric analysis.

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of developing International Standards is carried out through ISO technical committees. Every member body interested in a subject for which a technical committee has been authorized has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

International Standard ISO 2962 was developed by Technical Committee ISO/TC 34, *Agricultural food products*, and was circulated to the member bodies in February 1983.

It has been approved by the member bodies of the following countries:

| | | |
|---------------------|------------------------|-----------------------|
| Australia | Iran | Romania |
| Austria | Iraq | South Africa, Rep. of |
| Belgium | Kenya | Switzerland |
| Bulgaria | Korea, Dem. P. Rep. of | Tanzania |
| Canada | Korea, Rep. of | Turkey |
| Czechoslovakia | Malaysia | United Kingdom |
| Egypt, Arab Rep. of | Mexico | USSR |
| France | Netherlands | Venezuela |
| Germany, F. R. | New Zealand | Yugoslavia |
| Hungary | Poland | |
| India | Portugal | |

No member body expressed disapproval of the document.

This second edition cancels and replaces the first edition (i.e. ISO 2962-1974).

NOTE — The method specified in this International Standard has been developed jointly with the International Dairy Federation (IDF) and the Association of Official Analytical Chemists (AOAC) and will also be published by these organizations.

Cheese and processed cheese products — Determination of total phosphorus content — Molecular absorption spectrometric method

1 Scope and field of application

This International Standard specifies a molecular absorption spectrometric method for the determination of the total phosphorus content of cheese. The method is applicable to all kinds of cheese and to processed cheese products.

2 Reference

ISO 707, *Milk and milk products — Methods of sampling*.

3 Principle

Digestion of the cheese by sulfuric acid and hydrogen peroxide.

Formation of molybdenum blue by addition of molybdate-ascorbic acid solution. Molecular absorption spectrometric measurement, at a wavelength of 820 nm, of the blue colour formed.

NOTE — Dry ashing may be used, provided that the procedure gives the same results as the wet digestion.

4 Reagents

All reagents shall be of recognized analytical grade. The water used shall be distilled or deionized water, free from phosphorus compounds.

4.1 Concentrated sulfuric acid (H_2SO_4), $\rho_{20} = 1,84$ g/ml.

4.2 Hydrogen peroxide, solution containing about 30 g of H_2O_2 per 100 ml.

4.3 Molybdate-ascorbic acid solution.

4.3.1 Sodium molybdate solution.

Dissolve 12,5 g of sodium molybdate dihydrate ($\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$) in 5 mol/l sulfuric acid solution, dilute with this sulfuric acid solution to 500 ml, and mix.

4.3.2 Ascorbic acid, solution.

Dissolve 10 g of ascorbic acid ($\text{C}_6\text{H}_8\text{O}_6$) in water, dilute to 200 ml, and mix.

NOTE — This solution cannot be stored.

4.3.3 Mixed solution.

Immediately before use, mix 25 ml of solution 5.3.1 with 10 ml of solution 4.3.2, dilute to 100 ml with water, and mix.

4.4 Phosphorus, standard solution corresponding to 100 μg of P per ml.

Dry for at least 48 h about 1 g of potassium dihydrogenorthophosphate (KH_2PO_4) in a desiccator over an efficient desiccant, for example concentrated sulfuric acid.

Dissolve 0,439 4 g of the previously dried phosphate in water, dilute to 1 000 ml, and mix.

5 Apparatus

All glassware shall be thoroughly cleaned with a phosphorus-free detergent and rinsed with distilled water.

5.1 Analytical balance.

5.2 Device for grinding or grating cheese, capable of being easily cleaned.

5.3 Boiling water-bath.

5.4 Digestion flasks (Kjeldahl flasks or digestion tubes), of capacity 25 ml.

5.5 Heating apparatus.

5.5.1 Micro gas burners, for heating Kjeldahl flasks.

NOTE — Electric heaters may also be used.

ISO 2962-1984 (E)

5.5.2 Heating block, for heating digestion tubes.

5.6 Glass beads.

5.7 Graduated cylinders, of capacities 5 and 25 ml.

5.8 One-mark volumetric flasks, of capacities 50 and 100 ml.

5.9 One-mark pipettes, to deliver 1, 2, 3, 5 and 10 ml.

5.10 Spectrometer, suitable for measurements at a wavelength of 820 nm, equipped with cells of optical path length 10 mm.

6 Sampling

6.1 See ISO 707.

6.2 Store the sample in such a way that deterioration and change in composition are prevented.

7 Procedure

7.1 Preparation of the test sample

Remove the rind, smear or mouldy surface layer of cheese, in such a way as to provide a sample representative of the cheese as it is usually consumed. Grind or grate the sample by means of an appropriate device (5.2). Mix the ground or grated mass quickly, and if possible grind or grate a second time and again mix thoroughly. If the sample cannot be ground or grated, mix it thoroughly by intensive stirring and kneading.

Transfer the test sample to an airtight container to await analysis, which should be carried out as soon as possible after grinding or grating. If delay is unavoidable, take all precautions to ensure proper preservation of the sample and to prevent condensation of moisture on the inside surface of the container.

Clean the device after grinding or grating the sample.

7.2 Determination

7.2.1 Into a digestion flask (5.4) weigh, to the nearest 1 mg, a test portion of 0,5 to 1,0 g of the test sample.

Add three glass beads and 4 ml of the concentrated sulfuric acid (4.1).

NOTE — If the water content of the cheese is less than 50 % (*m/m*), a test portion of approximately 0,5 g is sufficient. In the case of fresh cheese, a test portion of approximately 1,0 g can be taken.

7.2.2 Heat the digestion flask under a well ventilated fume hood.

NOTE — A Kjeldahl flask shall be placed in an inclined position.

Control the height of the flame so as to limit the production of foam in the flask. Foaming into the neck of the flask is allowed but the foam shall not escape.

Keep the mixture gently boiling. Avoid local overheating and heating the flask above the surface of the liquid contents.

7.2.3 As soon as the foaming stops, cool to room temperature. Carefully add a few drops of the hydrogen peroxide solution (4.2) and reheat.

Repeat this procedure until the contents have become clear and colourless. During heating, mix the contents from time to time by careful swirling. Avoid local overheating.

7.2.4 Rinse the neck of the flask with about 2 ml of water. Heat the contents again until the water has evaporated.

Allow the liquid to boil for 30 min in order to destroy traces of hydrogen peroxide. Avoid local overheating.

7.2.5 Cool to room temperature. Quantitatively transfer the liquid contents into a 100 ml one-mark volumetric flask (5.8). Dilute to the mark with water and mix well.

7.2.6 Pipette 1 ml of the solution into a 50 ml one-mark volumetric flask (5.8) and dilute with about 25 ml of water. Add 20 ml of the molybdate-ascorbic acid solution (4.3.3). Dilute to the mark with water and mix well.

7.2.7 Heat the flask in the water-bath (5.3) for 15 min.

7.2.8 Cool to room temperature in a cold water-bath. Within 1 h, measure the absorbance of the solution against that of the blank test solution (see 7.4) at a wavelength of 820 nm.

7.3 Calibration curve

7.3.1 Pipette 10 ml of the standard phosphorus solution (4.4) into a 100 ml one-mark volumetric flask (5.8). Dilute to the mark with water and mix well.

7.3.2 Pipette into a series of five 50 ml one-mark volumetric flasks (5.8) 0, 1, 2, 3 and 5 ml, respectively, of the diluted standard solution (7.3.1), i.e. equivalent to 0, 10, 20, 30 and 50 µg of P respectively. Dilute the contents of each flask to approximately 20 ml with water.

7.3.3 Add to the contents of each volumetric flask 20 ml of the molybdate-ascorbic acid solution (4.3.3). Dilute with water to the mark and mix well.

Proceed as specified in 7.2.7.

7.3.4 Cool to room temperature in cold water. Within 1 h, measure the absorbance of each of the calibration solutions against water as reference at a wavelength of 820 nm.

7.3.5 Plot these absorbances against the amounts of phosphorus added.

7.4 Blank test

Carry out a blank test by following the procedure specified in 7.2, but without a test portion.

8 Expression of results

8.1 Method of calculation

The total phosphorus content, expressed as a percentage by mass, is equal to

$$\frac{m_1}{100 m_o}$$

where

m_o is the mass, in grams, of the test portion;

m_1 is the mass, in micrograms, of phosphorus, read from the calibration curve (or calculated from the regression line obtained by the method of least squares).

Report the result to the second decimal place.

8.2 Repeatability

The difference between two single results obtained on identical test material by one analyst using the same apparatus within a short time interval should exceed 0,03 g of phosphorus per 100 g of product on average not more than once in 20 cases in the normal and correct operation of the method.

8.3 Reproducibility

The difference between two single and independent results obtained by two operators working in different laboratories on identical test material should exceed 0,06 g of phosphorus per 100 g of product on average not more than once in 20 cases in the normal and correct operation of the method.

9 Test report

The test report shall show the method used and the result obtained. It shall also mention all operating conditions not specified in this International Standard, or regarded as optional, as well as any incidents that may have influenced the result.

The test report shall include all the information necessary for the complete identification of the sample.

STANDARD PREVIEW
 iTeh (standards.iteh.ai)

ISO 2962:1984

<https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984>

iTeh STANDARD PREVIEW
This page intentionally left blank
(standards.iteh.ai)

[ISO 2962:1984](https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984)

<https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984>

iTeh STANDARD PREVIEW
This page intentionally left blank
(standards.iteh.ai)

[ISO 2962:1984](https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984)

<https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984>

iTeh STANDARD PREVIEW
This page intentionally left blank
(standards.iteh.ai)

[ISO 2962:1984](#)

<https://standards.iteh.ai/catalog/standards/sist/a47396f2-7277-4ec8-8c78-eee5ac5a281f/iso-2962-1984>