

ISO/FDIS 30024:2023(E)

~~Date: 2023-09-25~~

ISO/TC 34/SC 10/~~AWG-12~~

Secretariat: INSO

Date: 2024-01-10

Animal feeding stuffs — Determination of phytase activity

Alimentation animale — Détermination de l'activité phytasique

iTeh Standards
(<https://standards.itih.ai>)

Document Preview
For FDIS stage

ISO/FDIS 30024

<https://standards.itih.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024>

© ISO 2024

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: + 41 22 749 01 11
E-mail: copyright@iso.org
Website: www.iso.org

Published in Switzerland

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[ISO/FDIS 30024](#)

<https://standards.iteh.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024>

Contents

Foreword	v
Introduction.....	vi
1 Scope	1
2 Normative references.....	1
3 Terms and definitions	1
4 Principle.....	3
5 Reagents.....	3
6 Apparatus.....	5
7 Sampling and sample preparation.....	6
8 Sample extractions	6
8.1 For compound feeds (excluding mineral feeds).....	6
8.2 For mineral feeds and premixtures.....	6
8.3 For feed additives	6
9 Procedure.....	7
9.1 General	7
9.2 Blank solution.....	7
9.3 Standards.....	7
9.3.1 Phosphate standard solution	7
9.3.2 Phytase level control.....	8
9.4 Standard curve.....	8
9.5 Phytase level control	9
9.6 For compound feed (excluding mineral feeds).....	9
9.7 For mineral feeds and premixtures.....	10
9.8 For feed additives	11
10 Calculations	12
10.1 Formulation of standard curve	12
10.2 Calculation of phytase activity.....	12
10.3 Correction for phytic acid purity and water content	13
10.4 Interference with blank values.....	14
11 Precision	14
11.1 Limit of detection and limit of quantification.....	14
11.2 Interlaboratory tests.....	15
11.3 Repeatability	15
11.4 Reproducibility	15
12 Test report.....	15
Annex A (informative) Interlaboratory study results for compound feeds (excluding mineral feeds) ..	16
A.1 First interlaboratory test.....	16

ISO/FDIS 30024:2024(E)

A.2	Second interlaboratory test.....	16
A.3	Third interlaboratory test on complementary compound feeds	18
Annex B (informative)	Interlaboratory study results for mineral feeds and premixtures.....	19
Annex C (informative)	Interlaboratory study results for feed additives	20
C.1	First interlaboratory test	20
C.2	Second interlaboratory test, specifically on 4a16 phytase feed additive	20
Bibliography	22

iTeh Standards (<https://standards.iteh.ai>) Document Preview

[ISO/FDIS 30024](https://standards.iteh.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024)

<https://standards.iteh.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024>

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO document should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

ISO draws attention to the possibility that the implementation of this document may involve the use of (a) patent(s). ISO takes no position concerning the evidence, validity or applicability of any claimed patent rights in respect thereof. As of the date of publication of this document, ISO had not received notice of (a) patent(s) which may be required to implement this document. However, implementers are cautioned that this may not represent the latest information, which may be obtained from the patent database available at www.iso.org/patents. ISO shall not be held responsible for identifying any or all such patent rights.

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 10, *Animal feeding stuffs*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 327, *Animal feeding stuffs - Methods of sampling and analysis*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

This second edition cancels and replaces the first edition (ISO 30024:2009), which has been technically revised.

The main changes are as follows:

- the scope has been extended to include complementary compound feeds, mineral feeds, premixtures and feed additives;
- phytic acid (phytate substrate) specifications have been added.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

This document quantifies phytase products in feeding stuff samples to control the phytase content of animal feed products. However, the method cannot be used to evaluate the *in vivo* efficacy of the phytase products.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[ISO/FDIS 30024](https://standards.iteh.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024)

<https://standards.iteh.ai/catalog/standards/iso/9dbfa537-9df3-45b6-b889-16128afb9359/iso-fdis-30024>

Animal feeding stuffs — Determination of phytase activity

1 Scope

This document specifies the determination of phytase activity in feeding stuff samples, including feed raw materials from plant origin, compound feeds (complete, complementary, mineral feeds), premixtures and feed additives.

The method is applicable to, and is collaboratively validated for, the determination of phytase activity in complete feed, complementary feed including mineral feed, premixtures and feed additives.

The method does not distinguish between phytase added as a feed additive and endogenous phytase already present in the feed materials. Therefore, the method is also applicable for feed materials from plant origin.

The method does not apply to evaluating or comparing the *in vivo* efficacy of the phytase product. It is not a predictive method of the *in vivo* efficacy of phytases present on the market as they can develop different *in vivo* efficacy per unit of activity.

2 Normative references

There are no normative references in this document.

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

— ISO Online browsing platform: available at <https://www.iso.org/obp>

— IEC Electropedia: available at <https://www.electropedia.org/>

3.1

phytase unit (U)

U

amount of enzyme that releases 1 μmol of inorganic phosphate from phytate per minute in acetate buffer at pH 5,5 and 37 °C

Note 1 to entry: This is determined by the reaction conditions specified in this document.

3.2

premixture

premix

uniform mixture of one or more micro-ingredients/*feed additives* (3.4) with a diluent and/or carrier, and that is not intended for direct feeding to animals

Note 1 to entry: Premixtures are used to facilitate the uniform dispersion of the micro-ingredients/additives in a larger mix.

[SOURCE: ISO 20588:2019, 3.2.39]

3.3

mineral feed

mineral mix

mineral supplement

feed that mainly consists of mineral elements, which is as an entire mix free-flowing

~~Note 1 to entry: In European legislation, a mineral feed contains at least 40 % of crude ash.~~

~~Note 2 to entry: to entry : Mineral feed is a form of *compound feed* (3.5) and of *complementary feed* (see Note 1 to entry of 3.5).~~

[SOURCE: ISO 20588:2019, 3.2.37, modified — "mineral feed" replaced "mineral mix" as the preferred term. Notes to entry added.]

~~Note 1 to entry: In European legislation, a mineral feed contains at least 40% of crude ash~~

~~Note 2 to entry : Mineral feed is a form of compound feed (3.5) and of complementary feed (note 1 of 3.5)~~

3.4 feed additive

substance intentionally added to feed and/or water, not consumed as feed by itself, whether or not it has a nutritional value, that affects the characteristics of feed including organoleptic properties, animal products, animal production or performance or welfare, or the environment

Note 1 to entry: Microorganisms, enzymes, acidity regulators, trace elements, vitamins and other products fall within the scope of this definition, depending on the purpose of use and the method of administration.

Note 2 to entry: Coccidiostats and histomonostats are a category of feed additives.

Note 3 to entry: Feed additive does not include *feed materials* (3.6) and *premixtures* (3.2).

[SOURCE: ISO 20588:2019, 3.2.18]

3.5 compound feed

formula feed
feed mixture

mixture of at least two *feed materials* (3.6), whether or not containing *feed additives* (3.4), for oral animal feeding in the form of a *complementary feed* or a *complete feed*

Note 1 to entry: *Complementary feed* is a form of *compound feed* as defined in ISO 20588:2019, 3.2.9.

Note 2 to entry: *Complete feed* is a form of *compound feed* as defined in ISO 20588:2019, 3.2.10.

[SOURCE: ISO 20588:2019, 3.2.11]

3.6 feed materials

products of vegetable or animal origin, whether or not containing *feed additives* (3.4), that are intended for use in oral animal feeding to meet animals' nutritional needs

Note 1 to entry: Feed materials can be in their natural state, fresh or preserved, or products derived from industrial processing, either organic or inorganic substances.

Note 2 to entry: Feed materials may be fed to animals either directly as such, or after processing, or in the preparation of *compound feed* (3.5), or as carrier of *premixtures* (3.2).

[SOURCE: ISO 20588:2019, 3.2.23, modified — "and products derived from industrial processing, either organic or inorganic substances" moved from the definition to Note 1 to entry.]

4 Principle

Phytase releases phosphate from the substrate myo-inositol hexakisphosphate (phytate). The released inorganic phosphate is determined by forming a yellow complex with an acidic molybdate/vanadate reagent. The optical density (OD) of the yellow complex is measured at a wavelength of 415 nm and the inorganic phosphate released is quantified from a phosphate standard calibration curve.

5 Reagents

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade and distilled or demineralized water or water of equivalent purity.

WARNING — This method requires the handling of hazardous substances. It is the responsibility of the user of this document to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

5.1 **Ammonia solution**, 25 % mass fraction; NH_3 .

5.2 **Ammonium heptamolybdate tetrahydrate**, $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot 4\text{H}_2\text{O}$.

5.3 **Ammonium monovanadate**, NH_4VO_3 .

5.4 **Hydrochloric acid**, 25 % mass fraction; HCl .

5.5 **Nitric acid**, 65 % mass fraction; HNO_3 .

5.6 **Potassium dihydrogenphosphate**, KH_2PO_4 .

5.7 **Phytate** (anionic form of phytic acid), **phytic acid**:

- all forms of phytate or phytic acid may be used, e.g. phytic acid (PA), phytic acid dodecasodium salt (Na-PA), phytic acid dodecapotassium salt (K-PA), phytic acid hexamagnesium salt n-hydrate (Mg-PA);
- $\leq 0,1$ % mass fraction of inorganic phosphorus;
- assay ≥ 90 % phosphorus (P) basis (dry basis);
- the substrate should have a percentage of IP6 (= hexaphosphate inositol containing six phosphate groups) of more than 95.

Information about IP6 ratio or percentage should be available from the supplier upon request.

As phytic acid salt hydrates are supplied with different contents of crystallization water, ensure that the crystallization water is in the stoichiometric range of 10 mol to 13 mol. In cases of deviation, see [10.3](#).

For control of the phytate, the blank OD from the standard curve (see [9.4](#)) shall be lower than 0,2. A higher OD value indicates phosphate or phytase contamination of used reagents.

5.8 **Sodium acetate trihydrate**, $\text{CH}_3\text{COONa}\cdot 3\text{H}_2\text{O}$.

5.9 **Polysorbate 20**¹.

¹ Tween® 20 is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of this product.

5.10 Diluted nitric acid.

Dilute one volume of nitric acid 65 % mass fraction (5.5) with two volumes water. Store at room temperature. The maximum storage time is two years.

5.11 Ammonium heptamolybdate reagent.

Dissolve 100,0 g of ammonium heptamolybdate tetrahydrate (5.2) in approximately 800 ml water (hot water at 50 °C to 60 °C may be used to facilitate salt dissolution). Add 10 ml 25 % mass fraction ammonia solution (5.1) and make up with water to 1 000 ml. Store at room temperature in the dark. The maximum storage time is two months.

5.12 Ammonium vanadate reagent.

Dissolve completely 2,35 g of ammonium monovanadate (5.3) in approximately 400 ml of water (hot water at 50 °C to 60 °C may be used to facilitate salt dissolution). Add 20 ml diluted nitric acid (5.10) and make up with water to 1 000 ml. Store at room temperature in the dark. The maximum storage time is two months.

5.13 Molybdate/vanadate STOP reagent.

Mix one volume of ammonium vanadate reagent (5.12) with one volume of ammonium heptamolybdate reagent (5.11) and add two volumes diluted nitric acid (5.10). Mix and store at room temperature. The maximum storage time is one day.

5.14 Polysorbate 20, 10 % mass fraction.

Dissolve 10,0 g of polysorbate 20 (5.9) with water and make up to 100 ml. Store at room temperature. The maximum storage time is six months.

5.15 Acetate buffer, pH 5,5; 0,25 mol/l.

Dissolve 34,0 g of sodium acetate trihydrate (5.8) in approximately 900 ml water. Adjust the pH with 25 % mass fraction hydrochloric acid (5.4) to $5,50 \pm 0,02$ and make up to 1 000 ml with water. Store at room temperature. The maximum storage time is two weeks.

5.16 Acetate buffer with 0,01 % mass fraction polysorbate 20, pH 5,5; 0,25 mol/l.

Dissolve 34,0 g of sodium acetate trihydrate (5.8) in approximately 900 ml water. Adjust the pH with 25 % mass fraction hydrochloric acid (5.4) to $5,50 \pm 0,02$. Add 1 ml 10 % mass fraction polysorbate 20 (5.14) and make up to 1 000 ml with water. Store at room temperature. The maximum storage time is two weeks.

5.17 Acetate buffer with 0,01 % mass fraction polysorbate 20, pH 5,5; 0,50 mol/l.

Dissolve 68,0 g of sodium acetate trihydrate (5.8) in approximately 900 ml water. Adjust the pH with 25 % mass fraction hydrochloric acid (5.4) to $5,50 \pm 0,02$. Add 1 ml 10 % mass fraction polysorbate 20 (5.14) and make up to 1 000 ml with water. Store at room temperature. The maximum storage time is two weeks.

5.18 Phytate substrate solution, 7,5 mmol/l (3 mmol/l end-concentration in the reaction).

Dissolve 2,00 g of phytate (5.7) in approximately 200 ml acetate buffer (5.15). Adjust the pH to $5,50 \pm 0,02$, e.g. with 25 % mass fraction hydrochloric acid (5.4), and make up with acetate buffer (5.15) to 250 ml. The maximum storage time is two weeks at 4 °C.

5.19 Phosphate stock standard solution, 50 mmol/l.