

# INTERNATIONAL STANDARD

ISO  
3100-2

First edition  
1988-07-15



INTERNATIONAL ORGANIZATION FOR STANDARDIZATION  
ORGANISATION INTERNATIONALE DE NORMALISATION  
МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ

## **Meat and meat products — Sampling and preparation of test samples —**

### **Part 2: Preparation of test samples for microbiological examination**

**iTeh Standards**  
**Document Preview**

*Viandes et produits à base de viande — Échantillonnage et préparation des échantillons pour essai —*

*Partie 2: Préparation des échantillons pour essai en vue de l'examen microbiologique*

[ISO 3100-2:1988](#)

<https://standards.iteh.ai/catalog/standards/iso/9ef6b82d-52e9-44e7-ae92-865a2a07ff01/iso-3100-2-1988>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council. They are approved in accordance with ISO procedures requiring at least 75 % approval by the member bodies voting.

<https://standards.iteh.ai>  
International Standard ISO 3100-2 was prepared by Technical Committee ISO/TC 34,  
*Agricultural food products.*

## Document Preview

ISO 3100 will consist of the following parts, under the general title *Meat and meat products – Sampling and preparation of test samples:*

[ISO 3100-2:1988](https://standards.iteh.ai/standard/iso/iso-3100-2-1988)

- [Part 1: Sampling](https://standards.iteh.ai/catalog/standards/iso/9ef6b82d-52e9-44e7-ae92-865a2a07ff01/iso-3100-2-1988)
- *Part 2: Preparation of test samples for microbiological examination*

# Meat and meat products — Sampling and preparation of test samples —

## Part 2: Preparation of test samples for microbiological examination

### 1 Scope

**1.1** This part of ISO 3100 gives general instructions and specifies procedures to be followed after taking a laboratory sample from meat and meat products, for the purpose of microbiological examination.

**1.2** A distinction is made between treatments for the following categories of products:

a) consignments or lots of meat or meat products prepared or packed as individual units of any size (for example sausages, vacuum-packed minced meat, sliced sausages, canned cooked ham), or meat in pieces not exceeding 2 kg in mass;

b) carcasses, cuts of carcasses, or cured meat in pieces exceeding 2 kg in mass, and mechanically separated meat or dried meat.

### 2 Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this part of ISO 3100. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this part of ISO 3100 are encouraged to investigate the possibility of applying the most recent editions of the standards listed below. Members of IEC and ISO maintain registers of currently valid International Standards.

ISO 3100-1 : —<sup>1)</sup>, *Meat and meat products — Sampling and preparation of test samples — Part 1: Sampling*.

ISO 6887 : 1983, *Microbiology — General guidance for the preparation of dilutions for microbiological examination*.

ISO 7218 : 1985, *Microbiology — General guidance for microbiological examinations*.

### 3 Principle

Preparation of test samples for microbiological examination. This may require the thawing and/or mincing of "open" meat samples or the pre-incubation, external sterilization, and aseptic opening of products processed or packed in sealed units.

### 4 Instructions of an administrative character

The sampling report and the label of the laboratory samples received should be checked (see ISO 3100-1). The date of receipt and the condition of the samples, including their temperature, should be noted. It should be clear what type of examination is to follow microbiological examination and whether the same or further samples will be used.

### 5 Diluents and reagents

#### 5.1 Base components

In order to improve the reproducibility of the results, it is recommended that, for the preparation of the diluents, dehydrated basic components should be used. Similarly, commercially prepared reagents may also be used. The manufacturer's instructions shall be rigorously followed.

The chemicals used shall be of analytical quality.

The water used shall be distilled or deionized, and shall be free from substances that might inhibit growth of micro-organisms under the test conditions.

pH measurements shall be carried out using a pH meter (6.10), adjusted to a temperature of 25 °C.

If the diluents and reagents are not used immediately, they shall, unless otherwise specified, be kept in the dark at a temperature between 0 °C and +5 °C, and in conditions that prevent any change in their composition. They shall not be kept for longer than 1 month.

1) To be published. (Revision of ISO 3100-1 : 1975.)

## 5.2 Diluent for cotton swabs.

### Composition

peptone	1,0 g
sodium chloride	8,5 g
water	1 000 ml

### Preparation and distribution of diluent

Dissolve the components in the water, by heating if necessary. Adjust the pH so that, after sterilization, it is 7,0 at 25 °C. Dispense into tubes or flasks of appropriate capacity in quantities such that, after sterilization, each tube or flask contains 9,0 ml of diluent.

Stopper the tubes or flasks.

Sterilize in the autoclave (6.1) at 121 °C ± 1 °C for 20 min.

## 5.3 Diluent for alginate swabs.

### Composition

sodium chloride	2,25 g
potassium chloride	0,105 g
calcium chloride	0,12 g
sodium hydrogen carbonate (NaHCO <sub>3</sub> )	0,05 g
sodium hexametaphosphate [mainly (NaPO <sub>3</sub> ) <sub>6</sub> ]	10 g
water	1 000 ml

### Preparation and distribution of diluent

Dissolve the components in the water, or dissolve commercially available tablets of the complete dry medium in 10 ml of water in tubes or flasks. If necessary, adjust the pH so that, after sterilization, it is 7,0 at 25 °C.

Dispense, if tablets have not been used, into stoppered tubes or flasks in such quantities that after sterilization each vessel contains 10 ml.

Sterilize in the autoclave (6.1) at 121 °C ± 1 °C for 20 min.

## 5.4 Ethanol, 95 % to 96 % (V/V).

## 5.5 Disinfectant mixture.

### Composition

ethanol (5,4)	60 ml
hydrochloric acid ( $\rho = 1,19$ g/ml)	10 ml
water	30 ml

## 6 Apparatus and glassware

NOTE — Disposable apparatus is an acceptable alternative to glassware if it has appropriate specifications.

Usual microbiological laboratory equipment and, in particular, the following.

## 6.1 Apparatus for dry sterilization (oven) or wet sterilization (autoclave).

Apparatus that will enter into contact with the diluents or the sample, except for apparatus that is supplied sterile (particularly plastics apparatus), shall be sterilized either

- by being kept at 170 °C to 175 °C for not less than 1 h in the oven (6.1), or
- by being kept at 121 °C ± 1 °C for not less than 20 min in the autoclave (6.1).

## 6.2 Blending equipment.

One of the following shall be used:

- a) mechanical meat mincer, laboratory size, capable of being sterilized, fitted with a plate with holes of diameter not exceeding 4 mm;
- b) peristaltic-type blender (Stomacher), with sterile plastic bags.

## 6.3 Incubators, for maintaining cans at a prescribed temperature for the detection of defective cans and for the rapid thawing of frozen samples.

## 6.4 Refrigerator, capable of being maintained at 2 °C, and a freezer capable of being maintained at or below –24 °C, for the storage of samples.

## 6.5 Instruments (sterilizable), for opening packages of meat and cutting up samples, for example can-openers, scissors, knives and forceps.

## 6.6 Swabs, made of cotton or alginate.

## 6.7 Tubes or flasks, with glass beads, in which swabs can be shaken.

## 6.8 Flasks, for drips from samples.

## 6.9 Pipettes or syringes, for removing drips from thawed or packaged meat samples.

## 6.10 pH meter, accurate to 0,1 pH unit at 25 °C.

## 7 Storage and receipt

### 7.1 General

Samples shall be stored at the prescribed temperature, protected from direct sunlight or other sources of heat.

Contamination shall be prevented (see also ISO 3100-1).

Start the examination as soon as possible after receipt of the samples, and in any case, within the limits given in 7.2 and 7.3.

## 7.2 Meat and meat products prepared or packed as individual units of any size, and meat in pieces not exceeding 2 kg in mass

### 7.2.1 Fresh meat

Store the samples in the refrigerator (6.4) on receipt and examine them within 24 h.