

---

INTERNATIONAL STANDARD



3199

---

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION · МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ · ORGANISATION INTERNATIONALE DE NORMALISATION

---

## Sodium chlorate for industrial use — Determination of chlorate content — Dichromate titrimetric method

*Chlorate de sodium à usage industriel — Dosage du chlorate — Méthode titrimétrique au dichromate*

First edition — 1975-02-15

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO 3199:1975](https://standards.iteh.ai/catalog/standards/sist/e56319bc-0ac3-439b-916a-672b16cf0ff4/iso-3199-1975)

<https://standards.iteh.ai/catalog/standards/sist/e56319bc-0ac3-439b-916a-672b16cf0ff4/iso-3199-1975>

---

UDC 661.442 : 546.135 : 543.24

Ref. No. ISO 3199-1975 (E)

**Descriptors** : sodium chlorate, chemical analysis, determination of content, chlorates, volumetric analysis.

## FOREWORD

ISO (the International Organization for Standardization) is a worldwide federation of national standards institutes (ISO Member Bodies). The work of developing International Standards is carried out through ISO Technical Committees. Every Member Body interested in a subject for which a Technical Committee has been set up has the right to be represented on that Committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the Technical Committees are circulated to the Member Bodies for approval before their acceptance as International Standards by the ISO Council.

International Standard ISO 3199 was drawn up by Technical Committee ISO/TC 47, *Chemistry*, and circulated to the Member Bodies in September 1973.

It has been approved by the Member Bodies of the following countries :

Austria	India	ISO 3199:1975
Belgium	Ireland	South Africa, Rep. of
Bulgaria	Italy	Spain
Chile	Netherlands	Switzerland
Czechoslovakia	New Zealand	Thailand
France	Poland	Turkey
Germany	Portugal	United Kingdom
Hungary	Romania	U.S.S.R.
		Yugoslavia

No Member Body expressed disapproval of the document.

# Sodium chlorate for industrial use – Determination of chlorate content – Dichromate titrimetric method

## 1 SCOPE AND FIELD OF APPLICATION

This International Standard specifies a dichromate titrimetric method for the determination of the chlorate content of sodium chlorate for industrial use.

The method is not applicable to the analysis of mixtures for agricultural use such as herbicides, pesticides, etc., in which certain constituents may cause interference.

## 2 PRINCIPLE

Reduction of the chlorate by a known quantity, in excess, of an iron(II) salt.

Titration of the excess of iron(II) salt with a standard volumetric solution of potassium dichromate in the presence of barium diphenylamine-4-sulphonate solution as indicator.

## 3 REAGENTS

During the analysis, use only reagents of recognized analytical reagent grade and only distilled water or water of equivalent purity.

**3.1 Orthophosphoric acid**,  $\rho$  approximately 1,71 g/ml, about 85 % (*m/m*) solution.

**3.2 Sulphuric acid**,  $\rho$  approximately 1,84 g/ml, about 96 % (*m/m*) solution or approximately 36 N.

**3.3 Sulphuric acid**, approximately 18 N solution.

**3.4 Iron(II) sulphate**, approximately 0,1 N solution.

Weigh, to the nearest 0,01 g, 39,2 g of ammonium iron(II) sulphate hexahydrate  $[(\text{NH}_4)_2\text{Fe}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}]$ , dissolve in 500 ml of water, slowly add 40 ml of the sulphuric acid solution (3.3), cool, dilute to the mark in a 1 000 ml one-mark volumetric flask and mix.

Store this solution in a glass bottle fitted with a ground glass stopper.

**3.5 Potassium dichromate**, 0,1 N standard volumetric solution.

Weigh, to the nearest 0,000 1 g, 4,903 g of potassium dichromate, previously dried at about 200 °C for about 16 h

and cooled in a desiccator. Dissolve in water, dilute to the mark in a 1 000 ml one-mark volumetric flask and mix.

Store this solution in a glass bottle fitted with a ground glass stopper.

**3.6 Barium diphenylamine-4-sulphonate**, 4 g/l solution.

## 4 APPARATUS

Ordinary laboratory apparatus.

## 5 PROCEDURE

### 5.1 WARNING

Sodium chlorate induces combustion. Avoid storing or handling it near a source of heat. Also avoid all contact of the product or its solutions with acids or combustible materials (paper, cardboard, clothing, wood, rags, fatty matter, etc.) which may ignite or subsequently form an explosive mixture. If any materials are accidentally impregnated with sodium chlorate, flush them with water.

### 5.2 Test portion

Weigh, to the nearest 0,001 g, approximately 3 g of the test sample.

### 5.3 Blank test

Carry out at the same time as the determination, following the same procedure, a blank test using the same quantities of all the reagents as used for the determination but replacing the test solution (5.4) by 20 ml of water.

### 5.4 Preparation of test solution

Dissolve the test portion (5.2) in water, dilute to the mark in a 1 000 ml one-mark volumetric flask and mix.

### 5.5 Determination

Place 20,0 ml of the test solution (5.4) in a 500 ml conical flask. Add 50,0 ml of the iron(II) sulphate solution (3.4), then add slowly and carefully, with cooling, 20 ml of the sulphuric acid solution (3.2) and 5 ml of the orthophosphoric acid solution (3.1). Allow to stand for 10 min at ambient temperature.

Dilute to approximately 300 ml, add 5 drops of the barium diphenylamine-4-sulphonate solution (3.6) and titrate with the standard volumetric potassium dichromate solution (3.5) until the colour changes to violet.

**6 EXPRESSION OF RESULTS**

**6.1 Method of calculation**

The sodium chlorate (NaClO<sub>3</sub>) content, expressed as a percentage by mass, is given by the formula

$$(V_1 - V_2) \times T \times \frac{1\,000}{20} \times \frac{100}{m} \times 0,017\,75 = \frac{88,75 (V_1 - V_2) T}{m}$$

where

*V*<sub>1</sub> is the volume, in millilitres, of the standard volumetric potassium dichromate solution (3.5) used for the blank test;

*V*<sub>2</sub> is the volume, in millilitres, of the standard volumetric potassium dichromate solution (3.5) used for the determination;

*T* is the actual concentration, expressed as normality, of the standard volumetric potassium dichromate solution (3.5);

*m* is the mass, in grams, of the test portion (5.2);

0,017 75 is the mass, in grams, of sodium chlorate corresponding to 1 ml of exactly 1 N standard volumetric potassium dichromate solution.

**6.2 Repeatability and reproducibility**

Comparative analyses carried out in 9 laboratories by 18 operators gave the statistical information shown in the following table :

<b>Mean (% m/m)</b>		99,55
<b>Standard deviation</b>	of repeatability ( $\sigma_r$ )	0,15
	of reproducibility ( $\sigma_R$ )	0,31

**7 TEST REPORT**

The test report shall include the following particulars :

- a) the reference of the method used;
- b) the results and the method of expression used;
- c) any unusual features noted during the determination;
- d) any operation not included in this International Standard, or regarded as optional.

<https://standards.iteh.ai/catalog/standards/sist/e56319bc-0ac3-439b-916a-672b16cf0ff4/iso-3199-1975>  
 ISO 3199:1975