



Designation: E1517 – 13

Standard Test Method for Determining the Effectiveness of Liquid, Gel, Cream, or Shampoo Insecticides Against Human Louse Ova¹

This standard is issued under the fixed designation E1517; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This test method determines the effectiveness of ovicidal materials in liquid, gel, cream, or shampoo form against the ova (that is, eggs or nits) of the human louse, *Pediculus humanus*.

1.2 This test method is intended for use by those wishing to develop efficacy data or compare ovicidal formulations for human louse ova control.

1.3 This test method consists of five replicates for a statistical comparison of formulations.

1.4 The values stated in SI units are to be regarded as standard. No other units of measurement are included in this standard, except for temperature measurements in Section 5.

1.5 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.*

2. Terminology

2.1 *Definitions of Terms Specific to This Standard:*

2.1.1 *hatched*—those eggs (nits) from which the nymph has emerged completely; an empty, clear egg case with the operculum clearly open.

2.1.2 *unhatched*—those eggs that are opaque; the operculum is closed or the nymph is partly emerged.

3. Summary of Test Method

3.1 Five replicates of 30 eggs are immersed in a test compound for a set period of time, washed, rinsed, blotted dry, and incubated.

3.2 Five control replicates are attached to human hair and processed as the treatment replicates, but with immersion in water.

3.3 Percent egg mortality, corrected by Abbott's Formula, is determined.

4. Significance and Use

4.1 This test method is a standardized test for the gathering of efficacy data for human louse ovicides.

4.2 Data collection in this manner is suitable for product development and comparison, and for review by regulatory agencies, to support the registration of human louse ovicidal products.

5. Apparatus and Materials

5.1 *Applicators*—Egg-infested hairs are attached to the end of a wooden applicator stick with duct tape such that 30 nits are on 1 to 3 hairs. Each replicate of 30 eggs is examined under a dissecting microscope to confirm viability. Any eggs that are shrunk or with other indications of being nonviable are excluded.

5.2 *Beakers*—A 100-mL beaker is used to contain 60 mL of test ovicide and another to contain 60 mL of water (control), into which the applicators are submerged. A 1000-mL beaker is used for washing the eggs.

5.3 *Heating Surface*—A slide dryer that provides heat of approximately 32°C (90°F).

5.4 *Incubator*, capable of maintaining a temperature of 31.7 ± 0.5°C (89°F) and a relative humidity of 60 ± 10 %.

5.5 *Water Bath*, capable of maintaining a temperature of 32°C (90°F).

5.6 *Wash Bottle, Stop Watch, and Dissecting Scope.*

5.7 *Test Insect*—The human louse, *Pediculus humanus*.²

5.8 *Positive Control Treatment (Optional)*—60 mL of solution known to give 65 to 95 % mortality of louse eggs when used under these test conditions.

¹ This test method is under the jurisdiction of ASTM Committee E35 on Pesticides, Antimicrobials, and Alternative Control Agents and is the direct responsibility of Subcommittee E35.12 on Insect Control Agents.

Current edition approved April 1, 2013. Published April 2013. Originally approved in 1993. Last previous edition approved in 2006 as E1517 – 99 (2006). DOI: 10.1520/E1517-13.

² A strain of the human body louse, *Pediculus humanus*, is maintained by Insect Control and Research, Inc., Baltimore, MD 21228-1199. The strain was established from a U.S. Department of Agriculture Gainesville colony. It is a susceptible strain and, through selection, has been adapted to the New Zealand White rabbit.