

FINAL
DRAFT

PUBLICLY
AVAILABLE
SPECIFICATION

ISO/DPAS
18999

ISO/TC 304

Secretariat: ANSI

Voting begins on:
2023-10-23

Voting terminates on:
2023-12-18

**Healthcare organization
management — Pandemic response
— Guidelines for respiratory infection
prevention and control in hospitals**

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[ISO/DPAS 18999](#)

<https://standards.iteh.ai/catalog/standards/sist/eef1861d-27bf-44bd-a422-36c3ba4b17db/iso-dpas-18999>

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.



Reference number
ISO/DPAS 18999:2023(E)

© ISO 2023

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

ISO/DPAS 18999

<https://standards.iteh.ai/catalog/standards/sist/eef1861d-27bf-44bd-a422-36c3ba4b17db/iso-dpas-18999>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2023

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

Page

Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions.....	1
4 Separate operation of a ward dedicated to respiratory infectious diseases.....	2
4.1 General.....	2
4.2 Principles of isolation room assignment and operation.....	2
5 Transportation of confirmed cases and roles of the dedicated healthcare team in a ward dedicated to respiratory infectious diseases.....	4
5.1 Patient transportation.....	4
5.1.1 Safe patient transportation.....	4
5.1.2 Roles and composition of dedicated patient transportation team.....	4
5.1.3 Precautions for patient transportation.....	5
5.1.4 Patient transportation ^[8]	6
5.1.5 Detailed handling procedures ^[9]	6
5.2 Roles of the dedicated healthcare team in a ward dedicated to respiratory infectious diseases and management of visitors (or guests).....	7
5.2.1 Roles and composition of the dedicated healthcare team in a ward dedicated to respiratory infectious diseases.....	7
5.2.2 Management of confirmed cases and visitors (or guests) in a ward dedicated to respiratory infectious diseases.....	8
6 Cleaning, disinfection and waste management.....	8
6.1 Disinfection of contaminated areas (rooms and equipment).....	8
6.1.1 Precautions for disinfection ^[10]	8
6.1.2 Disinfection methods.....	9
6.1.3 Precautions after disinfection ^{[7][6]}	11
6.2 Waste collection.....	12
6.2.1 Key measures for preventing the cross-infection of waste collectors.....	12
6.2.2 Principles for medical waste disposal.....	13
6.2.3 Wearing PPE during waste disposal.....	13
6.2.4 Safe waste disposal.....	13
Annex A (informative) Facility standards for isolation rooms in a ward dedicated to respiratory infectious diseases.....	15
Annex B (informative) Ward facility management for preventing cross-infection.....	18
Annex C (informative) Terminal cleaning checklist.....	26
Bibliography.....	28

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO document should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

ISO draws attention to the possibility that the implementation of this document may involve the use of (a) patent(s). ISO takes no position concerning the evidence, validity or applicability of any claimed patent rights in respect thereof. As of the date of publication of this document, ISO had not received notice of (a) patent(s) which may be required to implement this document. However, implementers are cautioned that this may not represent the latest information, which may be obtained from the patent database available at www.iso.org/patents. ISO shall not be held responsible for identifying any or all such patent rights.

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by ISO/TC 304, *Healthcare organization management*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

In the wake of the COVID-19 pandemic, guidelines have become necessary to prevent cross-contamination of the respiratory tract that can occur in hospitals in a disaster situation where a respiratory infectious disease has occurred. Therefore, this document was written to prevent cross-contamination in hospitals due to the outbreak of common respiratory infectious diseases.

This document is intended to standardize the guidelines for the separate operation of wards dedicated to respiratory infectious diseases; transportation of confirmed cases; and cleaning, disinfection and waste management, etc. in order to prevent respiratory cross-infections under emergency situations caused by respiratory infectious diseases such as COVID-19.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[ISO/DPAS 18999](https://standards.iteh.ai/catalog/standards/sist/eef1861d-27bf-44bd-a422-36c3ba4b17db/iso-dpas-18999)

<https://standards.iteh.ai/catalog/standards/sist/eef1861d-27bf-44bd-a422-36c3ba4b17db/iso-dpas-18999>

Healthcare organization management — Pandemic response — Guidelines for respiratory infection prevention and control in hospitals

1 Scope

This document provides guidelines to prevent cross-infections within a hospital, with a specific focus on the separate operation of wards dedicated to highly contagious respiratory infectious diseases, transportation of confirmed cases of highly contagious respiratory infectious diseases, disinfection, waste management, etc.

This document applies to the following:

- a) separate operation of wards dedicated to highly contagious respiratory infectious diseases;
- b) transportation of confirmed cases of highly contagious respiratory infectious diseases and roles of the dedicated healthcare team in a ward dedicated to highly contagious respiratory infectious diseases;
- c) cleaning, disinfection, and waste management.

2 Normative references

There are no normative references in this document.

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.1

coronavirus

virus that is part of a large family of viruses that can cause illness in animals or humans

Note 1 to entry: In humans, several coronaviruses are known to cause respiratory infections ranging from the common cold to more severe diseases such as Middle East Respiratory Syndrome (MERS) and Severe Acute Respiratory Syndrome (SARS). The coronavirus discovered in 2019 causes the coronavirus disease COVID-19.

[SOURCE: ISO 5472:2022, 3.3]

3.2

infectious disease patient

person who has tested positive in a diagnostic test for an infectious disease

3.3

negative pressure room

room in which the air pressure differential between the room and the adjacent indoor airspace directs the air flowing into the room (i.e. room air is prevented from leaking out of the room and into adjacent areas such as the corridor)

[SOURCE: ISO 5472:2022, 3.8]

3.4

internal corridor

corridor inside the negative pressure isolation area, which connects a patient room anteroom, corridor anteroom, personal protective equipment (PPE) *doffing room* (3.5), *waste disposal room* (3.6), *equipment storage room* (3.7), etc.

3.5

doffing room

space connected to the *internal corridor* (3.4), where healthcare workers who completed medical treatment take off their PPE before entering the general area

3.6

waste disposal room

space connected to the *internal corridor* (3.4), where medical waste generated from treatment of inpatients is sterilized or stored before discharge

Note 1 to entry: The waste disposal room is set to maintain a negative pressure lower than the internal corridor and a high temperature autoclave can be installed, if necessary.

3.7

equipment storage room

space connected to the *internal corridor* (3.4), where mobile equipment used for the treatment of *infectious disease patients* (3.2), etc. is stored or disinfected after use

3.8

HEPA filter

high efficiency particulate air filter retentive matrix having a minimum particle-collection efficiency of 99,97 % (that is, a maximum particle penetration of 0,03 % for 0,3 µm particles)

Note 1 to entry: See Reference [4].

[SOURCE: ISO 5472:2022, 3.6, "high efficiency particulate air filter" has been changed from a preferred term to an admitted term; note 1 to entry has been added.]

4 Separate operation of a ward dedicated to respiratory infectious diseases

4.1 General

A ward dedicated to respiratory infectious diseases refers to a unit that is established in order to prevent the infection of other patients and healthcare workers in the process of treating respiratory infectious disease patients, etc. and to suppress the transmission of causative pathogens in the local community. In principle, a negative pressure room should be operated as a single-occupancy room to curb the spread of infectious agents within a hospital through droplets or the air. See [Annex A](#).

4.2 Principles of isolation room assignment and operation

4.2.1

In principle, isolation rooms in a ward dedicated to respiratory infectious diseases should be operated as a single-occupancy negative pressure room, with the aim to prevent the circulation of air from isolation rooms occupied by confirmed cases to other areas within a hospital.

- a) When no negative pressure room is available, it is necessary to prevent the circulation of air from isolation rooms occupied by confirmed cases to other areas within the hospital as best as possible.

The criteria for air handling units (AHU) and heating, ventilation and air-conditioning (HVAC) system are as follows (see [Annex B](#)): because the default is to circulate a mix of outside air (30 %) and inside air (70 %), the opening rate of AHU is adjusted to prevent air mixing and to switch to a system with 100 % outside air supply and 100 % exhaust. In case of resource-limiting settings, staff open the windows at least three times a day for more than 30 min.

- b) When no single-occupancy room is available, multi-patient rooms can be used for confirmed cases in a ward that is completely separated from the routes of general patients.
- c) There are partial guidelines for the prevention and management of infectious diseases, so overall management guidelines are needed.

4.2.2

Suspected cases are assigned single-occupancy negative pressure rooms in principle until testing results are released. However, when no single-occupancy negative pressure room is available, patients with confirmed cases can be admitted to separate single-occupancy rooms that meet the AHU criteria.

4.2.3

The priority for the assignment of negative pressure rooms is to firstly assign them to high-risk patients in need of medical treatment, etc. The high-risk groups in terms of the priority for room assignment are as follows:

- a) patients with oxygen saturation below 90 % who require initial oxygen therapy;
- b) patients with underlying diseases (chronic obstructive pulmonary disease, cardiovascular diseases, etc.).

EXAMPLE Room assignment for confirmed cases in healthcare facilities^[5].

- 1) Priority is given to positive patients that are persons under investigation (PUI) or undergoing aerosol generating procedures (AGP).
- 2) Each confirmed case should be assigned to a single-occupancy negative pressure room in principle.
- 3) When no single-occupancy negative pressure room is available, a confirmed case should be assigned to a multi-patient negative pressure room.
- 4) When no multi-patient negative pressure room is available, a confirmed case should be assigned to a regular single-occupancy room.
- 5) When no regular single-occupancy room is available, a confirmed case should be assigned to a regular multi-patient room (a minimum distance of 1 m is recommended between beds).
- 6) When no regular multi-patient room is available, confirmed cases should be assigned to all rooms on a single floor in the facility.

The conditions for 3), 4), and 5) are as follows:

route: in order to ensure completely separate routes between confirmed cases and general patients, a ward (or single floor) is operated independently when confirmed cases are admitted to regular rooms.

5 Transportation of confirmed cases and roles of the dedicated healthcare team in a ward dedicated to respiratory infectious diseases

5.1 Patient transportation

5.1.1 Safe patient transportation

To ensure safe patient transportation, the following should be considered.

- a) A patient transportation team should be organized with a minimum number of personnel when transporting a confirmed case of a respiratory infectious disease.
- b) An advance notice should be provided to the receiving healthcare facility before the arrival of a confirmed case of a respiratory infectious disease so that proper preparations are taken prior to patient arrival.
- c) During the transportation of a confirmed case, any contact personnel should wear PPE.
- d) An ambulance should be used that is prepared at the transportation location as a vehicle for transportation.
- e) A patient should be transported on a predetermined route of the shortest travel distance and time (using designated elevators and pathways reserved exclusively for infectious disease patients).
- f) When transporting a patient to a ward dedicated to respiratory infectious diseases, the route should be blocked to prevent contact with other patients or visitors.
- g) The wheelchair or transportation cart used for patient transportation should be left in the ward dedicated to respiratory infectious diseases.

Patient transportation equipment, etc. that are left in the ward dedicated to respiratory infectious diseases can be reused after being disinfected according to the infection control guidelines.

- h) After transportation, PPE should be removed and disposed of; hand hygiene should be performed thoroughly^[6].

5.1.2 Roles and composition of dedicated patient transportation team

The dedicated patient transportation team for confirmed cases of respiratory infectious diseases performs the roles outlined in [Table 1](#).

Table 1 — Roles of dedicated patient transportation team for confirmed cases of respiratory infectious diseases

Category	Roles	Department or person in charge
Medical support	— Designation, cancellation, and management of hospital access and restricted areas. — Installing signs or signboards for restricted areas	Support department
	— Disinfection of surrounding environment after patient transportation, such as designated elevators, rooms, transportation route, etc.	Environmental service
	— Wearing a N95 or its equivalent or higher-grade respirator and PPE (excluding access control personnel who have no direct contact with patients and maintains a 2 m distance).	Administration personnel in charge of access control to patient transportation route
	— Access control of general patients, guardians and hospital staff to the transportation route before transporting confirmed cases.	
— Access control to reserved designated elevators.		

Table 1 (continued)

Category	Roles	Department or person in charge
	— Setting up a restricted area according to the hospital policies and attaching and installing signboards if necessary.	Administration personnel in charge of guidance on patient transportation and provision of related support
	— Facility management for cancellation and reoperation of restricted facilities.	
	— Disinfection of surrounding of environment.	
	— Wearing PPE and taking over a patient from emergency medical services (EMS) personnel.	Patient transportation team (two persons per team)
	— In the case of a severe patient, a healthcare worker accompanies the patient during transportation.	
	— Transporting a patient to an assigned room using transportation equipment (negative pressure stretcher, negative pressure wheelchair, etc.) depending on the patient's condition.	

5.1.3 Precautions for patient transportation

Matters related to patient transportation are at the discretion of healthcare workers; and these guidelines outline specific precautions for transportation.

a) Preparations for patient transportation

- 1) An ambulance should be used for patient transportation (an isolation stretcher equipped with a HEPA filter should be used, if available).
- 2) A patient transportation team should be organized with the minimum number of personnel (driver, health service workers, healthcare worker, etc.); it should be verified that there is no other person with the patient.
- 3) An advance notice should be provided to the receiving healthcare facility so that proper preparations are taken prior to patient arrival.
 - the following information should be delivered when requesting patient transfer, including:
 - patient condition (notable information such as severity, age, underlying diseases, dialysis status, cancer patient, mental illness, etc.);
 - patient location (name of the healthcare facility, etc.);
 - contact details of healthcare workers who can explain the patient's health condition.

b) Considerations for infection prevention^{[4][7]}

- 1) Patient transportation personnel should wear PPE during patient transportation according to [5.1.1](#).
- 2) Aerosol generating procedures should be prohibited whenever possible and aerosol-generating clinical pattern/procedures should be reduced before arriving at the hospital.
- 3) Any behaviour that can lead to pathogen transmission through contact during patient transportation should be avoided, such as taking off a mask, eating food, touching the face.
- 4) After patient transportation, the vehicle and transportation equipment should be cleaned and disinfected, given the possibility of pathogen transmission from the patient.

c) The patient should wear a mask during transportation, with the aim to minimize exposure through the respiratory system or physical contact; the patient should wear a mask whenever possible.