

SLOVENSKI STANDARD SIST EN ISO 11137-2:2012

01-oktober-2012

Nadomešča: SIST EN ISO 11137-2:2007 SIST EN ISO 11137-2:2007/AC:2009

Sterilizacija izdelkov za zdravstveno nego - Sevanje - 2. del: Določanje odmerka sterilizacije (ISO 11137-2:2012)

Sterilization of health care products - Radiation - Part 2: Establishing the sterilization dose (ISO 11137-2:2012)

iTeh STANDARD PREVIEW

Sterilisation von Produkten für die Gesundheitsfürsorge - Strahlen - Teil 2: Festlegung der Sterilisationsdosis (ISO 11137-2:2012)

SIST EN ISO 11137-2:2012

Stérilisation des produits de santé a/fradiation spartie 2. Établissement de la dose stérilisante (ISO 11137-2:2012)

Ta slovenski standard je istoveten z: EN ISO 11137-2:2012

<u>ICS:</u>

11.080.01 Sterilizacija in dezinfekcija na Sterilization and disinfection splošno in general

SIST EN ISO 11137-2:2012

en

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN ISO 11137-2:2012 https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012

SIST EN ISO 11137-2:2012

EUROPEAN STANDARD NORME EUROPÉENNE EUROPÄISCHE NORM

EN ISO 11137-2

March 2012

ICS 11.080.01

Supersedes EN ISO 11137-2:2007

English Version

Sterilization of health care products - Radiation - Part 2: Establishing the sterilization dose (ISO 11137-2:2012)

Stérilisation des produits de santé - Irradiation - Partie 2: Établissement de la dose stérilisante (ISO 11137-2:2012) Sterilisation von Produkten für die Gesundheitsfürsorge -Strahlen - Teil 2: Festlegung der Sterilisationsdosis (ISO 11137-2:2012)

This European Standard was approved by CEN on 14 March 2012.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.

https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

EN ISO 11137-2:2012 (E)

Contents	
----------	--

Annex ZA (informative) Relationship between this European Standard and the Essential Requirements of EU Directive 90/385/EEC on active implantable medical devices	4
Annex ZB (informative) Relationship between this European Standard and the Essential Requirements of EU Directive 93/42/EEC on medical devices	5
Annex ZC (informative) Relationship between this European Standard and the Essential Requirements of EU Directive 98/79/EC on <i>in vitro</i> diagnostic medical devices	6

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN ISO 11137-2:2012 https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012

Foreword

This document (EN ISO 11137-2:2012) has been prepared by Technical Committee ISO/TC 198 "Sterilization of health care products" in collaboration with Technical Committee CEN/TC 204 "Sterilization of medical devices" the secretariat of which is held by BSI.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by September 2012, and conflicting national standards shall be withdrawn at the latest by September 2012.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN ISO 11137-2:2007.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association, and supports essential requirements of EU Directive(s).

For relationship with EU Directive(s), see informative Annex ZA, B, C, which is an integral part of this document.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard; Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.^{37-2:2012}

https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178-

d461c274c0fa/sist-en-iso-11137-2-2012

Endorsement notice

The text of ISO 11137-2:2012 has been approved by CEN as a EN ISO 11137-2:2012 without any modification.

Annex ZA

(informative) Relationship between this European Standard and the Essential Requirements of EU Directive 90/385/EEC on active implantable medical devices

This European Standard has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association to provide a means of conforming to Essential Requirements of the New Approach Directive 90/385/EEC on active implantable medical devices.

Once this standard is cited in the Official Journal of the European Union under that Directive and has been implemented as a national standard in at least one Member State, compliance with the clauses of this standard given in Table ZA.1 confers, within the limits of the scope of this standard, a presumption of conformity with the corresponding Essential Requirements of that Directive and associated EFTA regulations.

Table ZA.1 — Correspondence be	etween this European Stand	ard and Directive 90/385/EEC
	incon this European otana	

Clauses of this International Standard	Essential Requirements (ERs) of EU Directive 90/385/EEC	Qualifying remarks/Notes
	<u>SIST EN ISO 11137-2:2012</u>	Standard and only in conjunction with ISO 11137-1. Packaging for maintenance of sterility during transportation and storage are not

d461c274c0fa/sist-en-iso-11137-2-2012

WARNING — Other requirements and other EU Directives may be applicable to the product(s) falling within the scope of this standard.

Annex ZB

(informative) Relationship between this European Standard and the Essential Requirements of EU Directive 93/42/EEC on medical devices

This European Standard has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association to provide a means of conforming to Essential Requirements of the New Approach Directive 93/42/EEC on medical devices.

Once this standard is cited in the Official Journal of the European Union under that Directive and has been implemented as a national standard in at least one Member State, compliance with the clauses of this standard given in Table ZB.1 confers, within the limits of the scope of this standard, a presumption of conformity with the corresponding Essential Requirements of that Directive and associated EFTA regulations.

Clauses of this Inte Standard	rnational	Essential Requirements (ERs) of EU Directive 93/42/EEC	Qualifying remarks/Notes
4, 5, 6, 7, 8, 9, 10		8.3 FANDARD PREVIE standards.iteh.ai) SIST EN ISO 11137-2:2012	This relevant ER is only partly addressed in this International Standard and only in conjunction with ISO 11137-1. Packaging for maintenance of sterility during transportation and storage are not covered.
4, 5, 6, 7, 8, 9, 10	+	n 8:4 catalog/standards/sist/14af1927-df70-4c 51c274c0fa/sist-en-iso-11137-2-2012	This relevant ER is addressed in this International Standard only in conjunction with ISO 11137-1.

Table ZB.1 — Correspondence between this European Standard and EU Directive 93/42/EEC

WARNING — Other requirements and other EU Directives may be applicable to the product(s) falling within the scope of this standard.

Annex ZC

(informative) Relationship between this European Standard and the Essential Requirements of EU Directive 98/79/EC on *in vitro* diagnostic medical devices

This European Standard has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association to provide a means of conforming to Essential Requirements of the New Approach Directive 98/79/EC on *in vitro* diagnostic medical devices.

Once this standard is cited in the Official Journal of the European Union- under that Directive and has been implemented as a national standard in at least one Member State, compliance with the clauses of this standard given in Table ZC.1 confers, within the limits of the scope of this standard, a presumption of conformity with the corresponding Essential Requirements of that Directive and associated EFTA regulations.

Clauses of this International Standard	Essential Requirements (ERs) of EU Directive 98/79/EC	Qualifying remarks/Notes
	<u>SIST EN ISO 11137-2:2012</u> tatdards.iteh.ai/catalog/standards/sist/14af192'	Standard and only in conjunction with ISO 11137-1. Packaging for maintenance of sterility during transportation and storage are not covered -8178-
4, 5, 6, 7, 8, 9, 10	B.2.4 ^{1461c274c0fa/sist-en-iso-11137-2-2}	This relevant ER is only addressed in this International Standard in conjunction with ISO 11137-1.

WARNING — Other requirements and other EU Directives may be applicable to the product(s) falling within the scope of this standard.

INTERNATIONAL STANDARD

Second edition 2012-03-15

Sterilization of health care products — Radiation —

Part 2: **Establishing the sterilization dose**

Stérilisation des produits de santé — Irradiation —

iTeh STPartie 2: Établissement de la dose stérilisante

(standards.iteh.ai)

SIST EN ISO 11137-2:2012 https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012



Reference number ISO 11137-2:2012(E)

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>SIST EN ISO 11137-2:2012</u> https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012



COPYRIGHT PROTECTED DOCUMENT

© ISO 2012

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office Case postale 56 • CH-1211 Geneva 20 Tel. + 41 22 749 01 11 Fax + 41 22 749 09 47 E-mail copyright@iso.org Web www.iso.org

Published in Switzerland

Contents

Forew	ord	v
Introd	uction	vi
1	Scope	1
2	Normative references	1
3 3.1 3.2	Terms, definitions and abbreviated terms Terms and definitions Abbreviated terms	1 1
4 4.1 4.2 4.3 4.4	Definition and maintenance of product families for dose setting, dose substantiation and sterilization dose auditing General Defining product families Designation of product to represent a product family for performance of a verification dose experiment or sterilization dose audit Maintaining product families	4 5 5
4.4	Effect of failure of establishment of sterilization dose or of a sterilization dose audit on a product family	
5 5.1 5.2 5.3 5.4 5.5	Selection and testing of product for establishing the sterilization dose	7 7 8 9 9
6	Methods of dose establishment	10
7 7.1 7.2 7.3 7.4	Method 1: dose setting using bioburden information Rationale Procedure for Method 1 for product with an average bioburden greater than or equal to 1,0 for multiple production batches Procedure for Method 1 for product with an average bioburden greater than or equal to 1,0 for a single production batch Procedure for Method 1 for product with an average bioburden in the range 0,1 to 0,9 for multiple or single production batches	11 12 17
8 8.1 8.2 8.3	Method 2: Dose setting using fraction positive information from incremental dosing to determine an extrapolation factor	20 21
9 9.1	Method VD _{max} — Substantiation of 25 kGy or 15 kGy as the sterilization dose Rationale	28 28
9.2	Procedure for Method VD_{max}^{25} for multiple production batches	
9.3 0.4	Procedure for Method VD_{max}^{25} for a single production batch	
9.4 9.5	Procedure for Method VD _{max} ¹⁵ for multiple production batches Procedure for Method VD _{max} ¹⁵ for a single production batch	
10 10.1	Sterilization dose audit Purpose and frequency	43

ISO 11137-2:2012(E)

10.2	Procedure for auditing a sterilization dose established using Method 1, Method 2A or Method 2B	43
10.3	Procedure for auditing a sterilization dose substantiated using Method VD_{max}^{25} or	
	Method VD _{max} ¹⁵	46
10.4	Failure of a sterilization dose audit	50
11	Worked examples	50
11.1	Worked examples Worked examples for Method 1	50
11.2	Worked examples for Method 2	53
11.3	Worked examples for Method VD _{max}	63
11.4	Worked example of a sterilization dose audit for a dose established using Method 1, the findings from which necessitated augmentation of the sterilization dose	
11.5	Worked example of a sterilization dose audit for a dose established using Method 2A, the findings from which necessitated augmentation of the sterilization dose	
11.6	Worked example of a sterilization dose audit for a sterilization dose substantiated using	
	Method VD _{max} ²⁵	67
Bibliog	Jraphy	69

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN ISO 11137-2:2012 https://standards.iteh.ai/catalog/standards/sist/14af1927-df70-4c8d-8178d461c274c0fa/sist-en-iso-11137-2-2012

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 11137-2 was prepared by Technical Committee ISO/TC 198, Sterilization of health care products.

This second edition cancels and replaces the first edition (ISO 11137-2:2006), which has been technically revised.

(standards.iteh.ai) ISO 11137 consists of the following parts, under the general title *Sterilization of health care products* — *Radiation*:

SIST EN ISO 11137-2:2012

- Part 1: Requirements for development, validation and routine control of a sterilization process for medical devices
- Part 2: Establishing the sterilization dose
- Part 3: Guidance on dosimetric aspects

Introduction

This part of ISO 11137 describes methods that can be used to establish the sterilization dose in accordance with one of the two approaches specified in 8.2 of ISO 11137-1:2006. The methods used in these approaches are:

- dose setting to obtain a product-specific dose;
- dose substantiation to verify a preselected dose of 25 kGy or 15 kGy.

The basis of the dose setting methods described in this part of ISO 11137 (Methods 1 and 2) owe much to the ideas first propounded by Tallentire^{[19][20][21]}. Subsequently, standardized protocols were developed^{[10][11]}, which formed the basis of the dose setting methods detailed in the AAMI Recommended Practice for Sterilization by Gamma Radiation^{[6][8]}.

Methods 1 and 2 and the associated sterilization dose audit procedures use data derived from the inactivation of the microbial population in its natural state on product. The methods are based on a probability model for the inactivation of microbial populations. The probability model, as applied to bioburden made up of a mixture of various microbial species, assumes that each such species has its own unique D_{10} value. In the model, the probability that an item will possess a surviving microorganism after exposure to a given dose of radiation is defined in terms of the initial number of microorganisms on the item prior to irradiation and the D_{10} values of the microorganisms. The methods involve performance of tests of sterility on product items that have received doses of radiation lower than the sterilization dose. The outcome of these tests is used to predict the dose needed to achieve a predetermined sterility assurance level (SAL).

Methods 1 and 2 can also be used to substantiate 25(kGy if, on) performing a dose setting exercise, the derived sterilization dose for an SAL of $10 T^{6}$ is less than or equal to 25 kGy. The basis of the method devised specifically for substantiation of 25 kGy, Method (VD_{max}) was put forward by Kowalski and Tallentire^[16]. Subsequent evaluations involving computational techniques demonstrated that the underlying principles were soundly based^[15] and field trials confirmed that Method VD_{max} is effective in substantiating 25 kGy for a wide variety of medical devices manufactured and assembled in different ways^[18].

A standardized procedure for the use of VD_{max} for substantiation of a sterilization dose of 25 kGy has been published in the AAMI Technical Information Report *Sterilization of health care products* — *Radiation sterilization* — *Substantiation of 25 kGy as a sterilization dose* — *Method* $VD_{max}^{[7]}$, a text on which the method described herein is largely based. Method VD_{max} is founded on dose setting Method 1 and, as such, it possesses the high level of conservativeness characteristic of Method 1. In a similar manner to the dose setting methods, it involves performance of tests of sterility on product items that have received a dose of radiation lower than the sterilization dose. The outcomes of these tests are used to substantiate that 25 kGy achieves an SAL of 10^{-6} .

To link the use of VD_{max} for the substantiation of a particular preselected sterilization dose, the numerical value of the latter, expressed in kilograys, is included as a superscript to the VD_{max} symbol. Thus, for substantiation of a sterilization dose of 25 kGy, the method is designated Method VD_{max}²⁵.

Method VD_{max}^{15} is based on the same principles as Method VD_{max}^{25} . The test procedure is similar to that of Method VD_{max}^{25} , but Method VD_{max}^{15} is limited to product with an average bioburden less than or equal to 1,5. The outcomes of the associated tests of sterility are used to substantiate that 15 kGy achieve a sterility assurance level of 10^{-6} .

This part of ISO 11137 also describes methods that can be used to carry out sterilization dose audits in accordance with ISO 11137-1:2006, Clause 12. Following establishment of the sterilization dose, sterilization dose audits are performed routinely to confirm that the sterilization dose continues to achieve the desired SAL.

Sterilization of health care products — Radiation —

Part 2: **Establishing the sterilization dose**

1 Scope

This part of ISO 11137 specifies methods for determining the minimum dose needed to achieve a specified requirement for sterility and methods to substantiate the use of 25 kGy or 15 kGy as the sterilization dose to achieve a sterility assurance level, SAL, of 10^{-6} . This part of ISO 11137 also specifies methods of sterilization dose audit used to demonstrate the continued effectiveness of the sterilization dose.

This part of ISO 11137 defines product families for sterilization dose establishment and sterilization dose audit.

2 Normative references il en STANDARD PREVIEW

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

SIST EN ISO 11137-2:2012

ISO 11137-1:2006, Sterilization of health care products 22 Radiation -81 Part 1: Requirements for the development, validation and routine control of a sterilization process for medical devices

ISO 11737-1, Sterilization of medical devices — Microbiological methods — Part 1: Determination of a population of microorganisms on products

ISO 11737-2, Sterilization of medical devices — Microbiological methods — Part 2: Tests of sterility performed in the definition, validation and maintenance of a sterilization process

3 Terms, definitions and abbreviated terms

For the purposes of this document, the terms and definitions given in ISO 11137-1 and the following apply.

3.1 Terms and definitions

3.1.1

batch

defined quantity of product, intended or purported to be uniform in character and quality, which has been produced during a defined cycle of manufacture

[ISO/TS 11139:2006, definition 2.1]

3.1.2

bioburden

population of viable microorganisms on or in product and/or sterile barrier system

[ISO/TS 11139:2006, definition 2.2]