
**Microbiology of the food chain —
Detection and quantification of
histamine in fish and fishery products
— HPLC method**

*Microbiologie de la chaîne alimentaire — Détection et quantification de
l'histamine dans le poisson et les produits de la pêche — Méthode CLHP*

Sample Document

get full document from standards.iteh.ai



Sample Document

get full document from standards.iteh.ai



COPYRIGHT PROTECTED DOCUMENT

© ISO 2017, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	1
5 Reagents and materials	1
6 Apparatus	2
7 Procedure	3
7.1 Sample preparation	3
7.2 Extraction	3
7.3 Derivatization.....	3
7.4 Purification.....	3
7.5 LC condition.....	4
7.6 Range of standard sample.....	4
8 Calculation	4
8.1 Calibration line (curve).....	4
8.2 Histamine quantification.....	5
9 Precision	5
9.1 Interlaboratory study.....	5
9.2 Repeatability limit.....	5
9.3 Reproducibility limit.....	5
Annex A (informative) Recommendations for HPLC separation	7
Annex B (informative) Performance characteristics of the method	9
Bibliography	14

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis — Horizontal methods*, in collaboration with ISO Technical Committee TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in accordance with the agreement on technical cooperation between ISO and CEN (Vienna Agreement).

Introduction

Histamine is a causative agent of scombroid poisoning or histamine fish poisoning. Histamine can be present mainly in *Scombridae* (tuna, mackerel) and *Clupeidae* (herring, sardine), species which contain a high level of free histidine. Histamine is formed through the decarboxylation of histidine by microbiological histidine decarboxylase.

Histamine [2-(1H-imidazol-5-yl)ethanamine] is defined as a biologically active low molecular weight basic nitrogenous molecule. The consumption of food containing significant concentration of histamine can cause symptoms similar to those associated to food allergies.

This document was developed in response to the need to standardize a method for histamine detection and quantification in fish and fishery products, in particular for European Regulation 2073/2005^[1] on microbiological criteria for foodstuffs.

Sample Document

get full document from standards.iteh.ai

Sample Document

get full document from standards.iteh.ai

Microbiology of the food chain — Detection and quantification of histamine in fish and fishery products — HPLC method

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices.

1 Scope

This document specifies a high performance liquid chromatography (HPLC) method to analyse histamine in fish and fishery products (fish sauces, fish matured by enzyme in brine, etc.) intended for human consumption.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

3 Terms and definitions

No terms and definitions are listed in this document.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

4 Principle

This method enables the separation of histamine among biogenic amines in fish and fishery products. The sample is extracted by mixing with perchloric acid. Precolumn derivatization is performed using dansyl chloride. The biogenic amines and the components in the solution are separated by HPLC with an appropriate column, using UV detection. Histamine mass concentration is calculated from the peak area ratio of histamine and internal standard with a calibration curve.

5 Reagents and materials

Use only reagents of recognized analytical grade and water complying with grade 1 of ISO 3696, unless otherwise specified. Solvents shall be of quality for HPLC analysis, unless otherwise specified.

5.1 Acetone, CH₃COCH₃.

5.2 Acetonitrile, CH₃CN.

5.3 Toluene, C₇H₈.