

FINAL DRAFT International Standard

ISO/FDIS 23691

Microbiology of the food chain — Determination and use of cardinal values

Microbiologie de la chaîne alimentaire — Détermination et utilisation des valeurs cardinales

nttps://standards.iteh Document Preview

ISO/TC 34/SC 9

Secretariat: **AFNOR**

Voting begins on: **2025-08-25**

Voting terminates on: 2025-10-20

CO/EDIC 22601

https://standards.iteh.ai/catalog/standards/iso/c83714ad-b998-44fb-9826-24d2d223b6c8/iso-fdis-23691

ISO/CEN PARALLEL PROCESSING

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

iTeh Standards (https://standards.iteh.ai) Document Preview

ISO/FDIS 23691

https://standards.iteh.ai/catalog/standards/iso/c83714ad-b998-44fb-9826-24d2d223b6c8/iso-fdis-23691



COPYRIGHT PROTECTED DOCUMENT

© ISO 2025

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office CP 401 • Ch. de Blandonnet 8 CH-1214 Vernier, Geneva Phone: +41 22 749 01 11 Email: copyright@iso.org Website: www.iso.org

Published in Switzerland

Contents				
Fore	eword		v	
Introduction				
1		oe		
_	-			
2		native references		
3	Terr	ns and definitions	2	
4	Principle			
	4.1	General		
	4.2	Gamma functions		
		4.2.1 General 4.2.2 Describing effects of temperature		
		4.2.2 Describing effects of temperature 4.2.3 Describing effects of <i>pH</i>		
		4.2.4 Describing effects of a_w		
		4.2.5 Describing the effects of concentrations of inhibitors		
	4.3	Process of cardinal values and food correction factor determination	10	
	4.4	Determination of the maximum specific growth rate		
		4.4.1 General	11	
		4.4.2 Binary dilution OD-based method		
		4.4.3 Direct plating method		
	4.5	Cardinal values determination		
	4.6 4.7	Validation		
	4.8	Use of cardinal values and food correction factor in predictions		
_				
5	Reag	gents and materials standards item. ai	15	
6	App	aratus	16	
7	Expo	erimental design and data collection	16	
	7.1	General		
	7.2	Preparation of culture and medium		
		7.2.1 Choice and storage of studied strain		
			22.17	
	7.2	7.2.3 Preparation of the modified nutrient brothLevels per factor to estimate cardinal values		
	7.3	7.3.1 General	_	
		7.3.2 Temperature		
		7.3.3 pH		
		7.3.4 Water activity		
		7.3.5 Inhibitory compounds		
	7.4	Experimental design to estimate the maximum specific growth rate from the binary dilution OD-based method		
	7.5	Experimental design to estimate the maximum specific growth rate from the direct plating method	23	
	7.6	Determination of the food correction factor based on a challenge test	23	
	7.7	Validation		
8	Evni	ession of the results: Estimation of the growth parameters	24	
J	8.1	General		
	8.2	Assessment of maximum specific growth rate at each level of intrinsic or extrinsic	1	
		factors (first step)	25	
		8.2.1 General		
		8.2.2 Assessment of maximum specific growth rates from direct plating data		
		8.2.3 Assessment of maximum specific growth rates by binary dilution OD-based		
		method	25	

	8.3	Assessment of cardinal values and optimum growth rate in broth, $\dot{\mu}_{\mathrm{Broth}}$ (second step)	
	8.3 8.4	Assessment of C_f (third step)	26 27
	9.1 9.2 9.3	General Prerequisites for growth predictions Using cardinal values to simulate growth 9.3.1 Growth simulation at a static given temperature 9.3.2 Growth prediction with dynamic time-temperature scenario	28 28 29 29
	Test r	9.3.3 Growth simulation at a static condition of temperature, pH and $a_{\rm w}$ eport	33
Annex	A (in	ty assurance if tools for primary and secondary fittings and ations	
		ormative) Guidance to obtain different $a_{ m w}$ values when using different humectants oth	37
Annex	C (info	ormative) Growth rate determination	38
Annex	D (info	ormative) Plate design	42
	variat	Formative) Example of the use of cardinal values for growth simulation and its tion	

<u>SO/FDIS 23691</u>

https://standards.iteh.ai/catalog/standards/iso/c83714ad-b998-44fb-9826-24d2d223b6c8/iso-fdis-23691

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO document should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

ISO draws attention to the possibility that the implementation of this document may involve the use of (a) patent(s). ISO takes no position concerning the evidence, validity or applicability of any claimed patent rights in respect thereof. As of the date of publication of this document, ISO had not received notice of (a) patent(s) which may be required to implement this document. However, implementers are cautioned that this may not represent the latest information, which may be obtained from the patent database available at www.iso.org/patents. ISO shall not be held responsible for identifying any or all such patent rights.

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 463, *Microbiology of the food chain*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

https://standards.iteh.ai/catalog/standards/iso/c83714ad-b998-44fb-9826-24d2d223b6c8/iso-fdis-23691

Introduction

Under the general principles of the Codex Alimentarius on food hygiene, it is the responsibility of the food business operators (FBOs) to control microbiological hazards in foods and to manage microbial risks. Therefore, it is the responsibility of the FBO to implement validated control measures, within the hazard analysis and critical control point (HACCP) system, and conduct studies in order to investigate compliance with the food safety criteria throughout the food chain.

In the framework of microbial risk assessment (MRA), several complementary approaches are developed to estimate risks posed by pathogens or spoilage microorganisms in the food chain. MRA is adopted by regulators under the auspices of the international agency for setting food standards. Predictive microbiology is one of the recognized scientific approaches used to validate control measures within the HACCP system, as well as to assess microbiological safety and quality of food, food production processes, food storage conditions and food preparation recommendations dedicated to consumers.

Therefore, this document provides technical rules, procedures and calculations to estimate the cardinal values of a microorganism of concern and use them in combination with challenge test results to simulate and predict its growth in raw materials, intermediate products or end products under reasonably foreseeable food processes, storage and use conditions.

To do so, this document includes the following sections:

- to identify the environmental factor(s) in scope (e.g. temperature, pH, a_w , organic acids);
- to define the appropriate experimental design;
- to estimate the cardinal values of a microorganism in broth medium;
- to perform a challenge test in the matrix of interest and derive the food correction factor and the maximum microbial population density;
- to use the cardinal values and the food correction factor to predict the growth of the studied microorganism in different conditions of interest (e.g. changes in time and temperature throughout the chill chain, changes in formulation with addition of organic acids or preservatives).

Regulatory authorities can have specific recommendations, and these differences have been included as much as possible in this document. It is, however, possible that additional requirements are needed to get a regulatory approval of the study.

The use of this document involves expertise from the organizing laboratories in relevant fields such as food microbiology, predictive microbiology and statistics. This expertise encompasses an understanding of sampling theory and design of experiments, statistical analysis of microbiological data, and overview of scientifically recognized and available mathematical concepts used in predictive microbiology.

Microbiology of the food chain — Determination and use of cardinal values

WARNING — In order to safeguard the health of laboratory personnel, it is essential that tests for detecting target microorganism(s) are only undertaken in properly equipped laboratories, under the control of a skilled microbiologist, and that great care is taken in the disposal of all incubated materials. Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety aspects, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices.

1 Scope

This document establishes basic principles and specifies requirements and methods to determine the cardinal values of bacteria and yeast strains and use them to predict microbial growth.

The four main steps of the approach are:

- a) determination of the cardinal values in culture medium;
- b) determination of the correction factor in the target food;
- c) validation of the model:
- d) simulations.

Four environmental factors are considered: temperature, pH, a_w and inhibitors (e.g. organic acids).

NOTE 1 Microbial competition is not considered as an inhibitor in this document and can be addressed by proper modelling approaches.

The determination of cardinal values is performed in a two-step approach: 24d2d223b6c8/iso-fdis-23601

- the determination of maximum specific growth rates of the studied strain grown in broth under a defined range of values of the studied environmental factor(s);
- the use of recognized predictive microbiology secondary models to fit the obtained experimental data to obtain the cardinal values.

The use of cardinal values in microbial growth simulation is based on predictive microbiology primary and secondary models. The cardinal values are combined with challenge test data to consider the matrix effect. Depending on the goal of the growth simulation, it is important to account for variation of cardinal values between strains within a bacterial or yeasts species.

Cardinal values are a good indicator of a strain growth ability for the studied environmental factors. They are therefore used as criteria to select strains, in addition to their origin and virulence, when performing growth challenge tests (see ISO 20976-1) or in methods validation (see ISO 16140 series).

NOTE 2 This document focuses on the determination of cardinal values for one strain. The same methodology can be used to characterize multiple strains independently to cover biological strain variability and include these results in the predictions.